

Pharmacotherapy Update: Fludarabine in the Treatment of Non-Hodgkin's Lymphoma

Lev Shvidel¹, Mirel Cohn² and Erica Sigler¹

¹Kaplan Medical Center, Rehovot, Israel. ²Assaf Harofeh Medical Center, Zerifin, Israel.

Abstract: Fludarabine, a purine nucleoside analogue, is considered as the most active drug in chronic lymphocytic leukemia. This review summarizes current knowledge concerning the mechanism of action, pharmacological properties, clinical activity and toxicity of fludarabine in the treatment of non-Hodgkin's lymphoma. A literature search on fludarabine for lymphomas was undertaken using MEDLINE databases. Clinical data shows that fludarabine alone and as a component of combination chemotherapy is effective in patients with various types of non-Hodgkin's lymphoma, particularly in follicular lymphoma. The most commonly used combinations are with cyclophosphamide and/or mitoxantrone. Fludarabine-based regimens are highly active in both previously untreated and relapsed indolent lymphomas. The results of the chemotherapy are considerably improved with the addition of rituximab. Fludarabine is generally well tolerated. Myelosuppression and infections, including opportunistic varieties, are the most frequent adverse effects. Indications for fludarabine and treatment regimens for lymphoma patients are discussed.

Keywords: fludarabine, Non-Hodgkin's lymphoma, treatment

Introduction

The success of the pyrimidine nucleoside cytarabine in the treatment of acute leukemias prompted development of new purine nucleoside analogues.¹ The most promising agents in this group are deoxycytosine (pentostatin), fludarabine phosphate and 2-chloro-2'-deoxyadenosine (2CdA) that cause suppression of DNA synthesis and repair in the tumor cells by distinct enzymatic processes. Fludarabine, synthesized by Montgomery and Hewson in 1969,² was first studied in acute leukemia using high doses (150 mg/m²/d for 5–7 days) and found to be excessively toxic to the nervous system.³ Subsequently fludarabine at lower doses revealed impressive activity in the treatment of indolent lymphoproliferative diseases.^{4–6} Currently fludarabine is considered as the most active drug in chronic lymphocytic leukemia (CLL) inducing complete remission (CR) in about 30% of previously untreated patients, with an overall response (OR) rate of 70%.^{7,8}

This review summarizes the available data on the fludarabine in the treatment of patients with non-Hodgkin's lymphomas (NHL). A literature search on fludarabine for lymphomas was undertaken to identify published guidelines, meta-analyses, systematic literature reviews and controlled clinical trials using MEDLINE databases. No evidence-based guidelines for the use of fludarabine have been found for the review. Pooling of data of cases from different studies for joint analysis was impractical for the reason that patients with heterogeneous characteristics (age, lymphoma subtype, prognostic factors, previous treatments etc) might have been included. The main goal of this study is to evaluate the majority of published records on fludarabine and to analyze the indications for the administration, the advantages and disadvantages of this drug in order to define its right current place in the treatment of lymphoma patients.

Mechanism of Action and Pharmacokinetics

Fludarabine, a fluorinated analog of adenine, is resistant to adenosine deaminase, the enzyme responsible for inactivation of the parent compound.⁹ In clinical practice it is used as a water soluble derivate, phosphate salt (9-β-D-arabinofuranosyl-2-fluoroadenine-5'phosphate). The pharmacokinetics of the

Correspondence: Lev Shvidel, MD, Hematology Institute, Kaplan Medical Center, Rehovot, Israel, 76100. Tel: (972) 89441383; Fax: (972) 89441706; Email: levsh@clalit.org.il



Copyright in this article, its metadata, and any supplementary data is held by its author or authors. It is published under the Creative Commons Attribution By licence. For further information go to: <http://creativecommons.org/licenses/by/3.0/>. The authors grant exclusive rights to all commercial reproduction and distribution to Libertas Academica. Commercial reproduction and distribution rights are reserved by Libertas Academica. No unauthorised commercial use permitted without express consent of Libertas Academica. Contact tom.hill@la-press.com for further information.

drug was studied after i.v. administration by rapid bolus injection and following short-term or continuous infusion. Fludarabine is rapidly dephosphorylated *in vivo* to F-ara-A showing peak plasma concentrations in a few minutes after the drug infusion. Following bolus injection, a brief 5 minutes early tissue-disposition phase is followed by 1–2 hours intermediate disposition phase and finally a prolonged terminal elimination phase (half-life 10 to 30 hours).¹⁰ After a prolonged intravenous infusion (>30 minutes) the disposition phase is less obvious, exhibiting a biphasic decline in plasma concentration of F-ara-A.^{11,12} No accumulation of plasma F-ara-A is seen following repeated daily administration. The main route of elimination is renal, 40%–60% of the fludarabine being excreted in the urine.¹³

The rephosphorylated product of F-ara-A, triphosphate F-ara-ATP, accumulates intracellularly within several hours (peak concentration at 4 hours) proportional to the dose administered, and has cytotoxic activity.¹⁴ The retention of F-ara-ATP varies between individuals, the half-life ranges from a few hours to several days with a median value of 15 hours.¹⁴ Fludarabine inhibits both DNA and RNA synthesis in the tumor cells by several mechanisms: 1. termination of chain elongation after incorporation into DNA strands, 2. inhibition of DNA and RNA polymerases, 3. inhibition of ribonucleotide reductase, the enzyme responsible for the conversion of ribonucleotides into deoxyribonucleotides, 4. direct activation of the apoptotic mechanisms.^{15,16} All the mentioned above data regarding the drug pharmacokinetics correlate with clinical results and should be applied in practical recommendations. The long half-life of the active metabolite allows for once-daily dosing. Patients with renal impairment and elderly patients who commonly have decreased creatinine clearance generally do not tolerate usual doses of fludarabine.¹⁷ Fludarabine dose reduction based on creatinine clearance values is needed, while the drug is relatively contraindicated in patients who have less than 30 ml/min.¹⁶ Finally, clinical results in CLL with divided doses administered on 3 or 5 consecutive days were preferable to the same accumulated dose given once weekly, indicating need to maintain therapeutic level for at least 3 days.¹⁸

Non-Hodgkin's Lymphomas

The NHLs comprise a diverse group of lymphoid malignancies with heterogeneous patterns of

molecular pathophysiology, histology, behavior and response to treatment. The modern classifications, like the World Health Organization (WHO) lymphoma classification, systematize lymphomas according to the characteristics of morphology, the origin and the state of differentiation of the tumor lymphocytes and specific cytogenetic aberrations.¹⁹ Using clinical criteria, the NHLs can be separated into two large groups: aggressive and indolent (low-grade). Aggressive lymphomas are associated with a rapidly progressive course, short survival if untreated, sensitivity to (immuno)chemotherapy and are treated with curative intent. The most common aggressive lymphoma is diffuse large B-cell lymphoma (DLBCL). Mantle cell lymphoma (MCL) is a distinct aggressive subtype showing low CR rate to therapy, median survival of only 3 to 4 years and few long-term survivors.

Indolent lymphomas constitute 30%–35% of all NHL subtypes. Of these, follicular lymphoma is the most encountered and diagnosed in the vast majority of patients. Follicular lymphoma has been associated with the chromosomal translocation t(14; 18), resulting in overexpression of the *bcl-2* oncogene and subsequent inhibition of apoptosis.

The less common subtypes are small lymphocytic lymphoma, nodal and mucosa-associated lymphoid tissue marginal zone lymphomas and lymphoplasmacytic lymphoma. Low grade lymphomas usually present with few symptoms, diffuse lymphadenopathy with or without splenomegaly and frequent bone marrow involvement and follow a long course. These lymphomas have tendency for good response to treatment but inevitable subsequent relapses. Low-grade lymphomas appear to be incurable with currently available treatments that rarely change the natural history, and death ultimately results from progression to more aggressive disease or infection. The overall median survival is 6 to 10 years.²⁰ The therapeutic approach to indolent lymphomas is controversial and ranges from “watch-and-wait” in asymptomatic patients or palliative chemotherapy to front-line myeloablative chemotherapy followed by autologous hematopoietic cells rescue in high-risk patients. In order to determine need for chemotherapy in follicular lymphoma, most clinicians use Follicular International Prognostic Index (FLIPI) score that assesses the risk of progression based on clinical stage, age, number of involved nodal sites, and hemoglobin and LDH levels (Table 1).²¹ The use of more active chemotherapeutic agents and novel

Table 1. Follicular lymphoma international prognostic index.²¹

1. age (>60 vs. ≤60)
2. Ann Arbor stage (III–IV vs. I–II)
3. hemoglobin level (<12 g/dL vs. ≥12 g/dL)
4. number of nodal areas (>4 vs. ≤4)
5. serum LDH level (>normal vs. normal)

Three risk groups are defined:

Low-risk (0–1 adverse factors, 36% of patients, 10-year survival of 70.7%);

Intermediate-risk (2 adverse factors, 37% of patients, 10-year survival of 51% and a hazard-ratio of 2.3);

Poor-risk (≥3 adverse factors, 27% of patients, 10-year survival of 35.5% and a hazard-ratio of 4.3).

therapeutic tools has recently produced complete responses and prolonged disease-free survival (DFS) intervals in most patients giving hope to survival curve improvement in indolent lymphomas.²² The means that have achieved the best results include monoclonal antibodies (including radioimmunotherapy) and purine analogs, namely, fludarabine.

Fludarabine Monotherapy

The initial study evaluating fludarabine in lymphoma was published by Leiby et al in 1987.²³ In this study a loading dose of 20 mg/m², followed by a continuous 48-hour infusion of 30 mg/m²/d, was administered to 26 patients with refractory disease. There were one CR which lasted for 9+ months and 7 short partial responses (PR) (OR 32%). Toxicity was acceptable and consisted mainly of myelosuppression.

Most patients enrolled in the early studies had advanced lymphoma refractory to previously given chemotherapy or progressed after response. Generally a schedule of fludarabine in a dosage of 20 to 30 mg/m²/day for 5 consecutive days and repeated every 3 to 5 weeks was applied. The responses, usually partial, have been achieved in approximately 50% of patients with indolent lymphomas and lasted up to one year (Table 2).^{4,24,25}

The results in aggressive NHL were disappointing showing low 0%–14% rate of objective responses.^{4,23}

The French cooperative group investigated fludarabine as first-line treatment in patients with advanced follicular lymphoma. Although the treatment was prematurely stopped because of toxicity

in nine patients, the response rate was relatively high: for 49 assessable patients the OR rate was 65% with CR rate of 37%; whereas the median time to progression was 13.6 months.²⁶ An Italian phase II trial used fludarabine monotherapy in 8 untreated and 13 previously treated stage 4 indolent lymphoma patients. The results were better in treatment-naïve (3 CR and 3 PR) than in pre-treated patients (0 CR and 8 PR).²⁷

The encouraging data demonstrated that fludarabine is capable to eradicate minimal residual disease (MRD) measured by polymerase chain reaction analysis (PCR) for bcl-2 in blood and bone marrow in some patients with follicular lymphoma.^{28–30} Although, there are controversies about its significance, the disappearance of MRD after therapy has been associated with significantly improved outcome.³¹ The MRD status may serve as good marker for remission quality, DFS and probably overall survival (OS).

Comparative trials

The Canadian investigators randomly compared the safety and efficacy of fludarabine and cyclophosphamide, vincristine, and prednisone (CVP-regimen) in patients with recurrent low-grade lymphoma after previous response to alkylator-based therapy.³² Although there were no significant differences in the response rates, the patients in the fludarabine arm had slightly improved median progression-free survival (PFS) and OS and better social function scores compared with patients treated with CVP. An OR rate of 64% (9% CR), a median PFS time of 11 months, and a median OS time of 57 months were demonstrated in the fludarabine group. Two patients died of fludarabine-related toxicities: invasive fungal infection and late development of Stevens-Johnson syndrome.

Recently the European Organization for Research and Treatment of Cancer (EORTC) Lymphoma Group reported long-term follow-up results of a phase III trial comparing fludarabine monotherapy to CVP-regimen in 381 previously untreated patients in advanced stages of indolent lymphoma. The responses rates were significantly improved in the fludarabine arm (70% OR and 38% CR vs. 52% and 15%, respectively, with CVP).³³ However, only non-significant trend for survival extension was noted: the 5-year OS in the “wait and watch” cohort treated by fludarabine was 65% while in the “intention to

Table 2. Noncomparative studies evaluating fludarabine monotherapy in non-Hodgkin's lymphoma.

Study	Patient numbers	Patients characteristics	OR %	CR %	Response duration, months	OS Median	Toxicity
Falkson ¹¹¹	21	pretreated indolent NHL	62	33	4.6	28+ months	moderate myelotoxicity
Fenchel et al. ¹¹²	14	pretreated indolent NHL	71	7	7	NR	infections, neutropenia
Foran et al. ⁷⁰	71	Untreated 35—LPC 19—WM 17—MCL	62 (in all) 63 79 41	15 (in all)	39 30 13	At 2.5 years not reached	myelotoxicity, rare infections
Gillis et al. ¹¹³	14	pretreated indolent NHL	51	1	NR	NR	rare infections
Hiddemann et al. ⁵	38	pretreated indolent NHL	31	13	12	NR	neutropenia
Hochster et al. ⁴	60	pretreated different NHL	30	15	NR	in CR/PR 30+ months	neutropenia, neurotoxicity
Leiby et al. ²³	26	pretreated different NHL	32	4	CR: 9+ PR 11 mo	NR	myelotoxicity
O'Brien et al. ¹¹⁴	48	Pretreated different NHL	50	10	14+	NR	myelotoxicity
Pigaditou et al. ²⁸	81	indolent NHL 45 pretreated 36 untreated	49	1	NR	NR	5 fatal infections, neutropenia
Redman et al. ²⁴	67	different pretreated lymphomas, (indolent, large-cell, MF, Hodgkin's)	37 (80 in follicular; 0 in aggressive NHL)	7	in CR 8–27+	NR	23 bacteremias, 2 PCP, 5 invasive fungal infection/fungemia
Solal-Céligny et al. ²⁶	49	Untreated follicular	65	37%	13.6	NR	neutropenia, few infections
Whelan et al. ²⁵	34	Pretreated indolent NHL	38	20	NR	NR	4 toxic deaths, infections
Zinzani et al. ²⁷	21	Follicular 8 untreated 13 pretreated	75	38	CR for 13+, 15+ and 17+	16 months	infections (1 fatal), neutropenia

Abbreviations: LPC, lymphoplasmacytic lymphoma; MF, mycosis fungoides.

treat" cohort was 76% (vs. 56% and 62%, respectively, in the CVP arm). Fludarabine-treated patients developed more commonly granulocytopenia, although it did not translate to more infections.

In contrast, the GELA study comparing fludarabine to CHVP (cyclophosphamide, doxorubicin, etoposide and prednisone) plus interferon in high-risk elderly patients with follicular lymphoma found better 2-year survival in the combination group.³⁴ Nevertheless, fludarabine in this study was given in much extended course (12 cycles within 2 years) that seems not optimal for elderly patients.

Combination (Immuno) Chemotherapy

The rationale for combining various chemotherapeutic agents with or without monoclonal antibodies is based on the different cytotoxic mechanisms of drugs, synergistic cytotoxicity in vitro and different adverse effects. An important additive cytotoxic activity of cyclophosphamide, anthracyclines or mitoxantrone and rituximab has been demonstrated when combined with fludarabine in vitro.^{35,36} Fludarabine inhibits repair of cyclophosphamide-induced cross links of the DNA.³⁷ Fludarabine also down-modulates the complement-resistance proteins CD55 and CD59, thereby potentially making cells more vulnerable to rituximab-induced complement-mediated lysis.³⁸ In addition fludarabine might (paradoxically) stimulate natural killer (NK) cell activity contributing to intensified tumor cell destruction by rituximab via antibody-dependent cell-mediated cytotoxicity (ADCC).³⁹ In turn, rituximab down-modulates IL-10 and bcl-2, making tumor lymphoid cells susceptible to fludarabine-induced apoptosis.⁴⁰ As a result, the combination of fludarabine and rituximab revealed cytotoxic activity on a lymphoma cell line resistant to either drug alone.³⁸

Combination with cyclophosphamide

Cyclophosphamide, the most frequently used alkylating agent in the treatment of lymphomas, was first described as therapy for diffuse large-cell lymphoma in 1975 and subsequently as a part of the CHOP regimen (cyclophosphamide, doxorubicin, oncovin, prednisone) became standard treatment for patients with aggressive lymphomas.^{41,42} CHOP-regimen induces very high (>90%) response rates in patients with indolent lymphoma and does not compromise subsequent stem cell

transplantation (SCT) feasibility.⁴³ However, the fact that this treatment is not able to change the natural history of indolent lymphomas or to eradicate MRD in follicular lymphomas, as documented by the persistence of cells bearing the bcl-2 gene rearrangement, has encouraged the investigation of cyclophosphamide in combination regimens with newer agents, such as fludarabine, mitoxantrone and rituximab.

The investigators from the Eastern Cooperative Oncology Group (ECOG) developed a regimen of cyclophosphamide and fludarabine (FC) (Table 3).⁴⁴ In this trial fludarabine (20 mg/m², days 1–5) was administered with an escalating dose of cyclophosphamide (600–1.000 mg/m², day1) in 27 previously untreated patients with indolent lymphomas. A 100% OR and 89% CR rates were achieved after a median number of six cycles. At a median follow-up of 61 months, 12 of these patients remain disease-free, and 5-year OS was 66%. Infectious complications including fungal sepsis, lobar pneumonia, *Pneumocystis carinii* pneumonia (PCP) (one of each) and three cases of *herpes zoster virus* (HZV) were reported.

Flinn et al from the John Hopkins Oncology Center using the FC regimen and myeloid growth factor support, achieved similar results in 20 untreated patients with follicular lymphoma: 92% OR rate with 60% CR.⁴⁵ A number of these patients were further autotransplanted; it is important to note that an autologous graft suitable for transplant was obtained in all patients seeking transplant.

Combination with mitoxantrone and anthracyclines

Mitoxantrone, an anthracenedione compound, has clinical activity similar to anthracyclines and is associated with less cardiac and gastrointestinal toxicity.⁴⁶ Response rates of 27%–67% have been reported in the management of indolent lymphomas, but generally the responses were partial.⁴⁶

A FND regimen, combined fludarabine (25 mg/m²/d on days 1–3), mitoxantrone (10 mg/m² on day 1) and dexamethasone (20 mg/d on days 1–5), was first described by McLaughlin et al from the M.D. Anderson Cancer Center (MDACC).⁴⁷ In this phase II trial, 51 patients with relapsed or refractory indolent lymphoma achieved an OR rate of 94% including 47% CR. The median duration of the CR was 21 months. Infections occurred in 12% of treatment cycles, including one fatal event

Table 3. Noncomparative studies evaluating fludarabine-based combination regimens in lymphoma.

Study	Patient numbers	Patients characteristics	Regimen	OR %	CR %	PFS median	OS median	Toxicity
Bordonaro et al. ¹¹⁵	18	elderly, indolent NHL	FND	94%	72	2-year 52%	2-year 67%	neutropenia, thrombopenia
Crawley et al. ⁴⁹	44	follicular NHL 10 untreated 34 pretreated (including 18 chemoresistant)	FND	chemo-responsive 84; chemo-resistant 44	20 (MR)	17 months	in PR/CR 30+ months	neutropenia, thrombopenia
Czuczman et al. ⁶¹	40	indolent NHL untreated pretreated	FR	88	81	not reached at 40+ months	not reached at 40+ months	mild
Hochster et al. ⁴⁴	27	indolent NHL untreated	FC	100	89	5-year DFS 53%	5-year 66%	fungal sepsis, pneumonia, PCP
Cohen et al. ¹¹⁶	30	MCL (10 untreated)	FC	63 (in all)	30	FFS: 5 months 28 months	17 months not reached at 42 mo	neutropenia, thrombopenia
McLaughlin et al. ⁴⁷	51	indolent NHL pretreated	FND	94	47	FFS: in CR 21 mo; in PR 9 months	NR	6 HZV, 2 PCP
Montoto et al. ³⁹	120	untreated FL	FCM	94 (81 MR)	83	5-year 58%	5-year 89%	Infections
Rohadgi et al. ¹¹⁷	37	untreated FL	Fludara followed by CNOP	97	67	34 months	not reached at 50 months	rare
Sacchi et al. ⁶³	54	pretreated FL	FCR	90 (86 MR)	74	8-year 61%	8-year 78%	few infections (one fatal)
Velasquez et al. ⁵⁰	78	indolent NHL untreated	FM	94	44	4-year 38%	4-year 88%	rare

Table 4. Comparative trials evaluating fludarabine-containing therapy.

Study	Patient numbers	Patients characteristics	regimen	OR	CR	PFS	OS	Toxicity
Forstpointer et al. ⁶⁵	128	pretreated indolent NHL and MCL	FCM vs. FCM-R	58%*	13%*	10 mo*	2-year 53%*	neutropenia, few infections
Foussard et al. ⁵⁴	142	elderly patients, indolent NHL	FM (reduced doses) vs. CHVP	79%*	33%*	16 mo*	73%*	neutropenia, few infections, more HZV in FM arm
Hagenbeek et al. ³³	381	untreated FL; intention to treat (<i>it</i>) or wait and watch (<i>w/w</i>)	Fludara vs. CVP	81%*	49%*	6 mo*	not reached at 53 months	neutropenia, few infections, more HZV in FM arm
				64%*	17%*	19 mo*		
				70%*	38%*	5-year <i>it</i> 17% <i>w/w</i> 8%	5-year <i>it</i> 76% <i>w/w</i> 65%	neutropenia
				52%*	15%*	<i>it</i> 19% <i>w/w</i> 24%	<i>it</i> 62% <i>w/w</i> 56%	
Klasa et al. ³²	91	recurrent indolent NHL	Fludara vs. CVP	64%	9%	11 mo	57 mo	3 toxic deaths in Fludara arm
McLaughlin et al. ⁶⁰	149	Untreated indolent NHL	FND plus rituximab concurrent vs. sequential	52%	7%	9 mo	44 mo	
				100%	92%;	3-year FFS 77%	3-year 95%	Neutropenia
				95%	85%	64%	95%	
Tsimberidou et al. ⁵²	142	Untreated indolent NHL	FND vs. ATT	97%	67%	5-year FFS 41%	5-year 84%	ATT more toxic
				97%	59%	50%	82%	
Zinzani et al. ⁵¹	199	indolent and MCL	Fludara vs. Fludara plus idarubicin	84%	47%	at 19 months 62% 84%	at 42 months 72% 72%	rare
Zinzani et al. ⁶²	140	untreated follicular NHL	FM vs. CHOP with or without sequential rituximab	81%	39%	3-year 63% no difference	3-year 94% no difference	CHOP more toxic
				96%	68%* (49% MR) 42%* (29% MR)	no difference		
				98%				

*statistically significantly different.

caused by *Mycobacterium avium-intracellulare*, 6 cases of dermatomal HZV and two PCP.

Other study has found similar clinical results with FND in previously treated patients.⁴⁸ Crawley et al demonstrated that the FND regimen is capable to induce molecular remission even in chemoresistant follicular lymphoma.⁴⁹

The Southwest Oncology Group (SWOG) in the U.S. conducted a trial of fludarabine and mitoxantrone as initial treatment in 78 indolent (in majority, follicular) lymphoma patients. This regimen produced OR rate of 94% and CR of 44%. The 4-year PFS of 38% and OS of 88% were favorable comparing to SWOG's prior experience with CHOP and ProMACE/MOPP (prednisone, methotrexate, doxorubicin, cyclophosphamide, etoposide, mechlorethamine, vincristine and procarbazine) regimens.⁵⁰ The toxicity was mild and transient.

Italian investigators designed a comparative trial of fludarabine (25 mg/m² for 3 days) in combination with idarubicin (12 mg/m² day 1) vs. fludarabine alone (at same daily dosage for 5 days).⁵¹ Following 6 monthly cycles 199 untreated patients with indolent lymphoma or MCL were assessed. The combination chemotherapy resulted in improved duration of response, although the response and OS rates as well as the rate of complications (a few in both groups) were comparable. Patients with follicular lymphoma had better CR rate (60% vs. 27%) and longer duration of response in comparison to MCL patients.

The MDACC group compared FND chemotherapy with intensive alternating triple therapy (ATT) that included 4 cycles of each CHOP-Bleo (CHOP and bleomycin), ESHAP (etoposide, methylprednisolone, cytarabine and cisplatin) and NOPP (mitoxantrone, vincristine, prednisone, procarbazine) regimens in a study on untreated patients with indolent lymphoma; maintenance with interferon/dexamethasone was given for one year in both treatment arms. The OR, OS and PFS rates were similar for FND and ATT.⁵²

In order to decrease myelotoxicity, Rohadgi et al used sequential therapy with 3 cycles of fludarabine as a single agent followed by 6–8 cycles of CVP with mitoxantrone (CNOP). The treatment was administered to 37 previously untreated patients with advanced follicular lymphoma and produced OR rate of 97% and CR rate of 67%. At median follow-up of 50 months, more than a half of responders were free of progression. Nevertheless,

two patients died of treatment-related events (disseminated HZV and secondary leukemia).⁵³ With the same intention to reduce toxicities, Foussard et al investigated lower doses of fludarabine in combination with mitoxantrone as first-line treatment and compared the results with cyclophosphamide, doxorubicin, vindesine and prednisone (CHVP-regimen) in 142 elderly patients with indolent lymphoma.⁵⁴ The FM therapy resulted in superior remission rates (OR 81% and CR 49% vs. 64% and 17%, respectively) and median failure-free survival (FFS) (36 months vs. 19 months). No difference was found between the study arms in terms of OS, however the lack of differences might be explained by efficient salvage treatment that often included fludarabine after failure of the CHVP. Despite common neutropenia, few infections (but more cases of HZV) were diagnosed in the FM arm.

The concomitant use of fludarabine, cyclophosphamide and mitoxantrone (FCM) in previously untreated patients induced excellent results of 94% OR, 83% CR and 81% molecular response rates.³⁰

Rituximab

Rituximab, a chimeric mouse/human antibody directed against CD20, is an important therapeutic option for patients with B-cell lymphomas. In preclinical studies, rituximab lysed CD20⁺ cells via human-complement, ADCC and by inducing apoptosis.⁵⁵ Rituximab as a single agent has significant clinical activity in follicular lymphomas, producing a clinical response rate of 73% and a molecular remission rate of 62%.^{56,57} When rituximab was combined with CHOP, the objective responses have been improved to >90% and CR was achieved in 50%–60% of the patients.⁵⁸ This immunochemotherapy resulted in longer DFS times and according to recent data probably improves also OS rates in patients with indolent lymphomas when compared with those who receive chemotherapy alone.⁵⁹ An additional advantage of rituximab resides in its good tolerability.

Investigators from the MDACC prospectively studied the FND regimen with rituximab given concurrently or sequentially as initial therapy in patients with stage IV small lymphocytic or follicular lymphomas.⁶⁰ Responding patients further received interferon- α 2b as maintenance therapy for a year. At a median follow-up of 30 months, 92% patients receiving concurrent treatment

Table 5. The most used fludarabine-containing regimens.***Fludarabine and rituximab (FR)**Fludarabine** 25 mg/m²/day i.v. (or 40 mg/m²/day p.o.***); days 1–5Rituximab 375 mg/m²/kg i.v.; day 1******Fludarabine, cyclophosphamide and rituximab (FCR)**Fludarabine 25 mg/m²/day, i.v. (or 40 mg/m²/day p.o.); day 1–3Cyclophosphamide 250 mg/m²/day, i.v. or p.o.; day 1–3Rituximab 375 mg/m² i.v.; day 1**Fludarabine, mitoxantrone (novantrone), rituximab and dexamethasone (FND-R)**Fludarabine 25 mg/m²/day, i.v.; day 1–3Mitoxantrone 10 mg/m², day 1, i.v.

Dexamethasone 20 mg/day, i.v. or p.o., day 1–5

Rituximab 375 mg/m², i.v.; day 1**Fludarabine, cyclophosphamide, mitoxantrone (novantrone) and rituximab (FND-R)**Fludarabine 25 mg/m²/day, i.v., day 1–3Cyclophosphamide 250 mg/m²/day, i.v., day 1–3Mitoxantrone 8 mg/m², i.v., day 1Rituximab 375 mg/m², i.v., day 1*In all regimens:*

*repeated every 28 days.

**dose adjustment for patients with renal failure (Table 6).

***oral fludarabine round dose to nearest 10 mg.

****for the first cycle, rituximab is recommended to administer on a day before chemotherapy. If the total WBC > 30 × 10⁹/L rituximab could be omitted from the first cycle of treatment to avoid risk of massive cytokine release syndrome. Rituximab can be incorporated in subsequent cycles of treatment regardless of the level of the WBC count.

achieved CR and 89% molecular remission, whereas the results were inferior (85% and 68%, respectively) post the sequential schedule.⁶⁰ The 3-year FFS and OS were statistically not different (77% and 95% for concurrent vs. 64% and 95% for sequential therapy, respectively). Although there was a slightly higher incidence of grade 3–4 neutropenia with concurrent treatment, there were no more infections.

Another phase II study investigated the efficacy of fludarabine and rituximab combination (FR) in 40 treatment-naïve or relapsed patients with indolent lymphomas that were enrolled on intent to treat basis.⁶¹ High and similar response rates were seen in treatment-naïve (88% OR and 81% CR) and previously treated (90% OR and 77% CR) patients. The median duration of response was not reached at 40 months. Out of 16 patients who were bcl-2-positive in bone marrow aspiration samples, 14 (88%) converted to negative after therapy and showed stable molecular response on repeated PCR assays for up to 4 years.

Italian investigators performed a landmark study to randomly compare the efficacy of fludarabine-mitoxantrone (FM) and CHOP regimens.⁶² Patients who achieved partial response (PR) or complete clinical but not molecular response were further treated with rituximab. Of 140 assessable patients, the OR rates for FM (96%) and CHOP (98%) were similar, but the rate of CR was higher in the FM arm (68% vs. 42%; *p* = 0.003). The rate of molecular response was also higher with FM (47% vs. 29%; *p* = 0.03). After rituximab, the number of patients who had experienced both clinical and molecular CR was higher in the FM arm (71% vs. 51%; *p* = 0.01). In the overall study population, rituximab treatment led to significantly higher clinical CR (86% vs. 57%) and combined CR and molecular response rates (61% vs. 29%). There were no differences in the PFS and OS times; the estimated survival rate for the entire study population at 3 years was 94%. Hematological toxicities were manageable in both arms, with no cases of treatment-related death or severe infection. CHOP was

Table 6. Dose modifications in patients with renal dysfunction.

Creatinine clearance (mL/min)	Dose	Fludarabine actual dose	
		i.v.	p.o
>70	100%	25 mg/m ² /d × 5d	40 mg/m ² /d × 5d
30–70	50%	20 mg/m ² /d × 3d	32 mg/m ² /d × 3d
<30		Do not use!!	

For any patient with a serum creatinine above normal and for all patients above the age of 60 years, creatinine clearance should be measured or calculated using the following formula to determine the initial dose of fludarabine.

$$\text{Estimated creatinine clearance (mL/minute)} = \frac{(140 - \text{age}) \times (\text{mass in kg}) \times [0.85 \text{ if female}]}{72 \times \text{serum creatinine (in mg/dL)}}$$

associated with significantly higher rates of grade 3–4 vomiting, alopecia, peripheral neuropathy and constipation. The results of this trial showed the advantage of the FM regimen over CHOP chemotherapy in terms of clinical and molecular responses, better tolerability, and also the benefit from the addition of rituximab that probably erased differences in the outcome.

Another Italian multi-center study investigated FCR regimen that was found as the most promising treatment in CLL.^{63,64} Following 4 cycles of FCR, excellent rates of response (74% CR and 84% molecular remission) were obtained among 54 patients with recurrent follicular lymphoma. At 45 months of follow-up the median OS and DFS had not been reached.

The German Low Grade Lymphoma Study Group prospectively compared FCM plus rituximab (FCM-R) with FCM alone in 128 patients with relapsed or refractory follicular, lymphoplasmacytic or mantle cell lymphoma, mostly post CHOP therapy.⁶⁵ The addition of rituximab significantly improved response rates: 79% OR and 33% CR in FCM-R arm as compared with 58% OR and 13% CR, respectively, for FCM alone ($p = 0.01$), with similar advantage in a subgroup analysis of follicular lymphoma (OR 94% vs. 70%) and MCL (58% vs. 46%, respectively). The R-FCM was significantly superior concerning PFS ($p = 0.038$) and OS ($p = 0.003$). There were no differences in the side effects in both study arms. This was the first prospective randomized trial to demonstrate that combined immunochemotherapy in patients with relapsed indolent and mantle cell lymphomas is superior to chemotherapy alone, both in terms of response and survival rates. The further trial of the German Group revealed significant prolongation of response period by

rituximab given as maintenance therapy in patients with relapsed or refractory indolent lymphoma and MCL, either or not they were previously treated by rituximab.⁶⁶

Radioimmunochemotherapy

The radionuclide applied in the radioimmune therapy has additive cytotoxic effect. Better results are obtained in patients with low tumor burden and non-severe bone marrow involvement. Leonard et al provided tumor mass reduction by an abbreviated course (3 cycles of fludarabine), followed by radioimmunotherapy with unlabeled anti-CD20 monoclonal antibody tositumomab and finally radiolabeled anti-CD20 monoclonal antibody iodine 131 tositumomab.⁶⁷ In this study on 35 patients with untreated follicular lymphoma fludarabine alone induced 89% response and 9% CR, and the full regimen—100% response with 86% CR. Ten out of 13 cases with baseline bone marrow bcl-2-positivity demonstrated molecular remission. The estimated 5-year PFS rate was 60%.

Fludarabine in Different Lymphoma Subtypes

The response rates were generally higher in patients with follicular lymphoma than in those with other subtypes, with the exception of follicular large-cell lymphoma (histological grade 3) or transformed follicular lymphomas. The results with fludarabine were worst in the diffuse large-cell lymphoma.⁹ MCL is characterized by especially low CR rate to therapy and bad prognosis.⁶⁸ Fludarabine monotherapy demonstrated limited efficacy in MCL inducing 33% PR rate with short (4 to 8 months) responses in a study on 15 patients, in majority pre-treated.⁶⁹ In previously untreated patients the

results were more encouraging (OR rates of 40%–70% with 30% CR),^{50,70} nevertheless, still not satisfactory. A number of small studies have investigated the efficacy of fludarabine-combined regimens in patients with MCL, usually as a subset of the entire cohort. The combination with mitoxantrone, idarubicin or cyclophosphamide produced OR rate of 60%–80% and CR rate of 20%–40% in either untreated or recurrent MCL patients.^{44,46,50} Although rituximab as a single agent in MCL revealed limited efficacy, in the setting of immunochemotherapy the results were significantly better compared to chemotherapy alone.⁶⁴

Several reports were dedicated to fludarabine therapy in lymphoplasmacytic lymphoma and its counterpart, Waldenstrom macroglobulinemia (WM), a clinicopathologic syndrome characterized by monoclonal immunoglobulin IgM production. Fludarabine salvage therapy induced response in a third of patients resistant to previously administered treatments.^{71–73} Despite a low rate of qualitative remissions, fludarabine provided useful palliation in this incurable disease. Furthermore, in a comparative trial on 90 patients with first relapse or with primary refractory WM, fludarabine was superior to cyclophosphamide-doxorubicin-prednisone (CAP) regarding response rates (30% vs. 11% PR) and duration (19 months vs. 3 months).⁷⁴ Fludarabine has been further studied as front-line treatment of WM: a European multicenter phase II study on 20 patients showed OR in 79%, with a median PFS of 40 months.⁷⁰ However, another large SWOG trial on 182 patients with symptomatic or progressive WM showed lower response rate (36% with only 2% CR). A serum beta2m level of ≥ 3 mg/L was the most significant adverse prognostic factor for survival in this study.⁷⁵

The success of fludarabine combinations in other lymphomas prompted similar studies also in WM.^{76,77} The best response rate of 95% was achieved by fludarabine and rituximab in 43 untreated or with < 2 prior therapies patients.⁷⁷ The median PFS was 51 months for all patients and 77 months for those previously untreated. However, relatively high rate of short term toxicities (6 pneumonias including 2 fatal) and long-term complications (3 cases of transformation to aggressive lymphoma and 3 cases of MDS/AML) should be evaluated in further studies.

High response rates (OR 91% and CR 53%) were achieved in marginal zone lymphoma in a

Greek trial evaluating FM combination followed by rituximab consolidation. At 39 months of median follow-up, 74% patients were alive and disease-free.⁷⁸

Finally, successful treatments with fludarabine in T-cell lymphomas, including mycosis fungoides, peripheral T cell lymphoma and angioimmunoblastic lymphadenopathy were occasionally reported.^{79–82}

Toxicity

Myelotoxicity and infection

The main toxic events in fludarabine treatment are associated with myelosuppression manifested by transient granulocytopenia, and immunosuppression related to profound decrease in lymphocytes.⁸³ The depletion in CD4 cells may persist for more than a year, while other lymphocyte populations recover more rapidly. The risk for infectious complications increases during the periods of treatment-related granulocytopenia, and the wide spectrum of infections includes bacterial and various opportunistic organisms. The most encountered are *Streptococcus pneumoniae*, *Staphylococcus* and *Haemophilus influenzae*, whereas gram-negative bacteria are common reason for hospitalization. Early studies with fludarabine have been associated with very high rate of complicated granulocytopenia and infections. For example, in 76 patients with relapsed lymphoma Redman et al diagnosed 23 episodes of fever of unknown origin, 23 cases of bacteremia, 3 of systemic candidiasis, 2 of PCP, one of each *aspergillosis* pneumonia, *Torulopsis glabrata* infection and disseminated *cytomegalovirus*, and 8 patients with localized HZV.²³ Poor performance status, elderly age, advanced or refractory lymphoma, concurrent corticosteroids and prolonged leukopenia predispose to infections. Restriction of concurrent corticosteroids, appropriate prophylactic use of cotrimoxazole and in special cases also acyclovir and antifungal drugs, prophylaxis by myeloid growth factors, doses reduction for elderly patients prevent many infectious events. However, these recommendations are mostly empiric and may be excessive for low-risk patients. PCP prophylaxis might not be required at least in younger patients receiving first-line chemotherapy.⁸⁴

Finally, cases of hepatitis B virus reactivation and acute hepatitis were diagnosed following fludarabine therapy.⁸⁵ Patients who are positive for

HBsAg or HBcoreAg should be treated with lamivudine for the entire duration of chemotherapy and for 6 months afterwards and monitored with liver function tests and hepatitis B virus DNA.

Neurotoxicity

Recommended doses of fludarabine for treatment of lymphomas very rarely cause neurological effects. Nevertheless, paralysis, neuropathy, blurred vision, seizures or headache were occasionally reported.⁸⁶ Progressive multifocal leucoencephalopathy (PML) associated with John Cunningham virus was described in lymphoma patients treated by fludarabine.^{63,87} Elderly patients are more prone to neurotoxicity.

Impaired stem cell collection

Younger patients with indolent lymphoma commonly become candidates for stem cell autotransplantation. It was shown that fludarabine may impair collection of peripheral blood progenitor cells destined for autotransplant in acute leukemia and CLL.⁸⁸ This observation could complicate the decision to choose this drug for patients who may require autotransplant in the future. However, at least a part of unsuccessful stem cells mobilizations in the reported lymphoma cases was performed in unsatisfactory circumstances: either too late in the course of lymphoma, or too soon after fludarabine treatment and/ or with myeloid growth factors alone.⁸⁹ The data collected by European Blood and Marrow Transplantation (EBMT) CLL subcommittee on 41 patients who had previously been treated by fludarabine, was encouraging.⁹⁰ The numbers of CD34+ cells collected at least 2 months elapsed between the last dose of fludarabine and the start of mobilizing therapy, was satisfactory. More CD34+ cells were collected in the patients who were treated with fewer than six courses of fludarabine compared to ≥ 6 courses and in patients who had received less chemotherapy before the mobilization. All 16 patients who were autotransplanted, engrafted.

The collection of stem cell early enough in the course of disease, the delay in stem cell collection for at least 2–3 months after fludarabine administration and the combination of chemotherapy and myeloid growth factors for stem cell mobilization can improve the quality of stem cell harvest.

Secondary malignancy

There is data suggesting that fludarabine may increase the risk of secondary malignancies, particularly

when combined with other DNA-damaging agents, such as alkylators or topoisomerase II inhibitors.^{91–93} The estimated rate for MDS/secondary AML reach 3%–4% for lymphoma patients treated with fludarabine-based chemotherapy given as first-line treatment and up to 10% for previously heavily treated patients. The complex cytogenetic aberrations and abnormalities of chromosomes 5 and/or 7 were common. The clinicians should discuss this issue with their (particularly younger) patients before the treatment administration.

Other events

Fludarabine induces little significant nausea, vomiting and alopecia. Tumor lysis syndrome associated with fludarabine treatment has been documented, usually in CLL and occasionally in lymphoma patients, so prophylactic measures should include hydration and allopurinol.⁹⁴ Fludarabine has been implicated in the development or worsening of autoimmune hemolytic anemia (AIHA) and thrombocytopenia (ITP) in CLL patients.⁹⁵ The administration of fludarabine needs caution when hemolytic anemia or acute thrombocytopenia have been diagnosed. The combination of fludarabine and cyclophosphamide and/or rituximab appears to reduce the incidence of autoimmune cytopenias.⁹⁶ Finally, transfusion associated graft-versus-host-disease after fludarabine treatment has been noted, so the use of irradiated blood products for transfusion following treatment is mandatory.⁹⁷

Oral Fludarabine

The oral formulation of fludarabine phosphate has been available since early 2000s. The 10 mg tablet contains an immediate-release form of the drug. Pharmacokinetics studies demonstrated that a once-daily oral dose of 40 mg/m² corresponded to i.v. exposure of 25 mg/m².⁹⁸ F-ara-A reaches peak plasma concentration following an oral dose in a 1½ hour. The bioavailability of the oral drug ranges from 30% to 80% between patients, while the intra-individual variations in bioavailability are low and not affected by meal.⁹⁹

A phase II multi-center study was conducted in Japan to assess the efficacy of oral fludarabine in 46 patients with indolent lymphoma (41 follicular) and 6 with MCL, previously treated by chemotherapy with or without rituximab.¹⁰⁰ The OR rate in the cohort of the indolent lymphoma patients was 65% with 30% CR, the responders showed

median time to treatment failure of 12 months. Only one patient with MCL experienced an objective response, the others 5 had stable disease through the treatment period.

The comparison of oral fludarabine alone or in combination with cyclophosphamide to results in similar patients treated by intravenous administration, showed same clinical efficacy. Similar tolerability profile was also found, except more gastrointestinal adverse events for the oral formulation.^{101,102}

Fabbri et al used reduced dosages of oral fludarabine (40 mg/day) and oral cyclophosphamide (150 mg/m²/day) (for 4 days and 4 cycles) in untreated elderly patients with indolent lymphoma.¹⁰³ The OR rate was 84% (CR 40%) with median PFS of 20 months.

In conclusion, the oral formulation of fludarabine is effective and attractive for oncologists who do not need the hospital day care facilities and also for patients who benefit from home conditions.

An Italian trial used this drug together with mitoxantrone and radioimmunotherapy (yttrium-90 radiolabeled antibody zevalin) in untreated patients with advanced indolent lymphoma.¹⁰⁴ When treatment accomplished, 55 out of 57 patients achieved CR, an estimated 3-year PFS was 76% and OS 100%.

Discussion Regarding the Choice of Treatment for Lymphoma Patients

Clinical evidence from studies in patients with lymphoma suggests that fludarabine monotherapy is at least as effective, if not better, than other chemotherapy such as CVP and CHOP-regimens in follicular and other types of indolent lymphomas and carries acceptable toxicity rate. At the contrary, fludarabine is not suitable for large-cell lymphoma. In comparative studies, fludarabine combined with cyclophosphamide and/or mitoxantrone improved the remission rate and duration when compared with single-agent fludarabine in both previously treated and untreated patients with indolent lymphoma. Finally, it is currently agreed that immunochemotherapy approach produces superior results to chemotherapy alone, whereas rituximab serves as the most effective and safe monoclonal antibody for the combinations. The question that still remains unresolved, which is the most effective combination chemotherapy regimen to use along with rituximab? The main candidates

are CHOP and fludarabine-based regimens. Unfortunately, there is no proof that any of the known chemotherapy regimen determines prolonged overall survival. As a result, no established recommendations for treatment of choice exist in indolent lymphoma. In the near future the results of the ongoing European intergroup study PRIMA (Primary Rituximab and Maintenance) that evaluates differences in the efficacy between different upfront R-chemo regimens (R-CVP, R-CHOP, R-FCM), as well as the potential impact of rituximab maintenance may shed light on this issue.

Currently the U.S. Guidelines from the National Comprehensive Cancer Network (NCCN) recommend various options as the first-line treatment of follicular lymphoma including: CVP and CHOP-regimens, fludarabine as a single agent or fludarabine-based regimens, all of them combined with rituximab; radioimmunotherapy or rituximab as a single therapy.¹⁰⁵ Rituximab maintenance after first-line therapy is recommended for further investigation following favorable results in pretreated patients.^{66,106}

For the elderly patients rituximab alone or a single alkylator agent may be selected. For the subsequent treatments bendamustine, chemoimmunotherapy as described for initial therapy, radioimmunotherapy, autologous or allogeneic SCT in selected patients can be used. Fludarabine as a component of combination therapy is also a recommended option for first-line therapy in small lymphocytic and marginal cell lymphoma and as second-line in MCL.

Data from the National LymphoCare survey in the United States demonstrated that lack of specified recommendations weakens the tailoring of the most efficient treatment program. Currently only few patients with follicular lymphoma are treated on clinical trials, whereas the clinicians use diverse regimens based on their personal preference.¹⁰⁷ Approximately half (49%) of them utilized chemoimmunotherapy, the others—"wait and watch" strategy (19%), rituximab monotherapy (13%), chemotherapy alone (4%) or radiation alone (5%). The CHOP-R and CVP-R regimens were favored (78%) when immunochemotherapy was administered, while combination of fludarabine-based regimen with rituximab was selected in 11%. The survey showed also significant differences in physicians' preferences in various medical centers and regions suggesting absence of a universally recognized policy. For example, initial watch and wait

was used in 31% of patients in the Northeast but in 13% in the Southeast, whereas fludarabine was used in 18% of patients in the Southwest and only 3% in the Northeast. M. Czuczman entitled his article “Controversies in follicular lymphoma: “Who, what, when, where, and why?” (Not necessarily in that order!)”, and emphasized that achievement of complete hematological and molecular remission has been associated with improvement in outcome.¹⁰⁸ His main practical conclusion: “suboptimal therapy in = suboptimal results out”.

What are the therapy preferences in our Hematology center? The goal of therapy for the younger patients with follicular lymphoma, requiring chemotherapy, is to achieve the best qualitative response and thus to prevent as possible relapse of the disease. We generally choose combination of fludarabine, cyclophosphamide and rituximab (FCR) as one of the most active regimens that demonstrates the highest clinical and molecular response rates. In cases with massive organomegaly we usually choose CHOP-R regimen because of possibility of hidden transformation of the disease. As known fludarabine-containing

regimens are poorly effective in cases with histologic transformation as well as in follicular large-cell lymphoma. In these settings, anthracycline-containing combinations are superior.^{109,110} The first best response to chemotherapy we use for autologous stem cell harvest (Fig. 1), that can be utilized in a subsequent relapse. We believe that with cautions that were discussed above, successful stem cell harvesting can be performed in most patients in remission.

In the elderly patients in addition to the decrease of tumor mass, the main goal of treatment is the maintenance of a good quality of life, rather than complete eradication of clinical and laboratory evidence of disease. Factors influencing survival in elderly population, apart from lymphoma parameters and prior chemotherapy, include performance status, underlying comorbidities, socioeconomic status and availability of social support. Chlorambucil (with or without prednisone) as an oral, well-tolerated and inexpensive drug is a reasonable first-line option in fragile patients. Combination of rituximab with COP or fludarabine in lower doses, which appears to be still active and has fewer associated toxicities than used in standard dose,




Figure 1. The proposed management of patients with follicular lymphoma.

may be an alternative. We administer fludarabine in a half of the standard cycle dosage and discontinue therapy when a satisfactory clinical response is achieved.

In conclusion, immunochemotherapy based on fludarabine-including regimens exhibits a high efficacy in the treatment of indolent and particularly follicular lymphoma.

Authors' Declaration

1. The article is original, has not been published in any other journal, is not under consideration by any other journal and does not infringe any existing copyright or any other third party rights.
2. We are the sole authors of the article and have full authority to enter into this agreement and in granting rights to *Libertas Academica* are not in breach of any other obligation.
3. The article contains nothing that is unlawful.
4. The authors have no competing financial interests.

To our and currently accepted scientific knowledge all statements in the article are true.

References

1. Ellison RR, Holland JF, Weil M, et al. Arabinosyl cytosine: A useful agent in the treatment of acute leukemia in adults. *Blood*. 1968;32:507–23.
2. Montgomery JA, Hewson K. Nucleosides of 2-fluoroadenine. *Journal of Medicinal Chemistry*. 1969;12:498–504.
3. Warrell RP Jr, Berman E. Phase I and II study of fludarabine phosphate in leukemia: therapeutic efficacy with delayed central nervous system toxicity. *Journal of Clinical Oncology*. 1986;4:74–9.
4. Hochster HS, Kim KM, Green MD, et al. Activity of fludarabine in previously treated non-Hodgkin's low-grade lymphoma: results of an Eastern Cooperative Oncology Group study. *Journal of Clinical Oncology*. 1992;10:28–32.
5. Hiddemann W, Unterhalt M, Pott C, et al. Fludarabine single-agent therapy for relapsed low-grade non-Hodgkin's lymphomas: a phase II study of the German Low-Grade Non-Hodgkin's Lymphoma Study Group. *Seminars in Oncology*. 1993;20(5 Suppl 7):28–31.
6. Zinzani PL, Lauria F, Rondelli D, et al. Fludarabine: an active agent in the treatment of previously-treated and untreated low-grade non-Hodgkin's lymphoma. *Annals of Oncology*. 1993;4:575–8.
7. Keating MJ, O'Brien S, Lerner S, et al. Long-term follow-up of patients with chronic lymphocytic leukemia (CLL) receiving fludarabine regimens as initial therapy. *Blood*. 1998;92:1165–71.
8. Johnson S, Smith AG, Löffler H, et al. Multicentre prospective randomised trial of fludarabine versus cyclophosphamide, doxorubicin, and prednisone (CAP) for treatment of advanced-stage chronic lymphocytic leukaemia. The French Cooperative Group on CLL. *Lancet*. 1996;347:1432–8.
9. Anderson VR, Perry CM. Fludarabine: a review of its use in non-Hodgkin's lymphoma. *Drugs*. 2007;67:1633–55.
10. Malspeis L, Grever MR, Staibus AE, Young D. Pharmacokinetics of 2-F-ara-A (9-beta-D-arabinofuranosyl-2-fluoroadenine) in cancer patients during the phase I clinical investigation of fludarabine phosphate. *Seminars in Oncology*. 1990;17(5 Suppl 8):18–32.
11. Danhauser L, Plunkett W, Liliemark J, Gandhi V, Iacoboni S, Keating M. Comparison between the plasma and intracellular pharmacology of 1-beta-D-arabinofuranosylcytosine and 9-beta-D-arabinofuranosyl-2-fluoroadenine 5'-monophosphate in patients with relapsed leukemia. *Leukemia*. 1987;1:638–43.
12. Adkins JC, Peters DH, Markham A. Fludarabine. An update of its pharmacology and use in the treatment of haematological malignancies. *Drugs*. 1997;53:1005–37.
13. Lichtman SM, Etcubanas E, Budman DR, et al. The pharmacokinetics and pharmacodynamics of fludarabine phosphate in patients with renal impairment: a prospective dose adjustment study. *Cancer Investigation*. 2002;20:904–13.
14. Gandhi V, Kemena A, Keating MJ, Plunkett W. Cellular pharmacology of fludarabine triphosphate in chronic lymphocytic leukemia cells during fludarabine therapy. *Leukemia Lymphoma*. 1993;10:49–56.
15. Pettitt AR. Mechanism of action of purine analogues in chronic lymphocytic leukaemia. *British Journal of Haematology*. 2003;121:692–702.
16. Robertson LE, Chubb S, Meyn RE, et al. Induction of apoptotic cell death in chronic lymphocytic leukemia by 2-chloro-2'-deoxyadenosine and 9-beta-D-arabinosyl-2-fluoroadenine. *Blood*. 1993;81:143–50.
17. Shvidel L, Shtalrid M, Bairey O, et al. Israeli Study Group on CLL. Conventional dose fludarabine-based regimens are effective but have excessive toxicity in elderly patients with refractory chronic lymphocytic leukemia. *Leukemia Lymphoma*. 2003;44:1947–50.
18. Kemena A, O'Brien S, Kantarjian H, et al. Phase II clinical trial of fludarabine in chronic lymphocytic leukemia on a weekly low-dose schedule. *Leukemia Lymphoma*. 1993;10:187–93.
19. Harris NL, Jaffe ES, Diebold J, et al. The World Health Organization classification of neoplasms of the hematopoietic and lymphoid tissues: report of the Clinical Advisory Committee meeting—Airlie House, Virginia, November, 1997. *Hematology Journal*. 2000;1:53–66.
20. Portlock CS. Management of the indolent non-Hodgkin's lymphomas. *Seminars in Oncology*. 1980;7:292–301.
21. Solal-Céligny P, Roy P, Colombat P, et al. Follicular lymphoma international prognostic index. *Blood*. 2004;104:1258–65.
22. Liu Q, Fayad L, Cabanillas F, et al. Improvement of overall and failure-free survival in stage IV follicular lymphoma: 25 years of treatment experience at The University of Texas M.D. Anderson Cancer Center. *Journal of Clinical Oncology*. 2006;24:1582–9.
23. Leiby JM, Snider KM, Kraut EH, Metz EN, Malspeis L, Grever MR. Phase II trial of 9-beta-D-arabinofuranosyl-2-fluoroadenine 5'-monophosphate in non-Hodgkin's lymphoma: prospective comparison of response with deoxycytidine kinase activity. *Cancer Research*. 1987;47:2719–22.
24. Redman JR, Cabanillas F, Velasquez WS, et al. Phase II trial of fludarabine phosphate in lymphoma: an effective new agent in low-grade lymphoma. *Journal of Clinical Oncology*. 1992;10:790–4.
25. Whelan JS, Davis CL, Rule S, et al. Fludarabine phosphate for the treatment of low grade lymphoid malignancy. *British Journal of Cancer*. 1991;64:120–3.
26. Solal-Céligny P, Brice P, Brousse N, et al. Phase II trial of fludarabine monophosphate as first-line treatment in patients with advanced follicular lymphoma: a multicenter study by the Groupe d'Etude des Lymphomes de l'Adulte. *Journal of Clinical Oncology*. 1996;14:514–9.
27. Zinzani PL, Lauria F, Rondelli D, et al. Fludarabine: an active agent in the treatment of previously-treated and untreated low-grade non-Hodgkin's lymphoma. *Annals of Oncology*. 1993;4:575–8.
28. Pigaditou A, Rohatiner AZ, Whelan JS, et al. Fludarabine in low-grade lymphoma. *Seminars in Oncology*. 1993;20(5 Suppl 7):24–7.
29. Zinzani PL, Pulsoni A, Perrotti A, et al. Fludarabine plus mitoxantrone with and without rituximab versus CHOP with and without rituximab as front-line treatment for patients with follicular lymphoma. *Journal of Clinical Oncology*. 2004;22:2654–61.
30. Montoto S, Moreno C, Domingo-Doménech E, et al. High clinical and molecular response rates with fludarabine, cyclophosphamide and mitoxantrone in previously untreated patients with advanced stage follicular lymphoma. *Haematologica*. 2008;93:207–14.

31. López-Guillermo A, Cabanillas F, McLaughlin P, et al. The clinical significance of molecular response in indolent follicular lymphomas. *Blood*. 1998;91:2955–60.
32. Klasa RJ, Meyer RM, Shustik C, et al. Randomized phase III study of fludarabine phosphate versus cyclophosphamide, vincristine, and prednisone in patients with recurrent low-grade non-Hodgkin's lymphoma previously treated with an alkylating agent or alkylator-containing regimen. *Journal of Clinical Oncology*. 2002;20:4649–54.
33. Hagenbeek A, Eghbali H, Monfardini S, et al. *Journal of Clinical Oncology*. 2006;24:1590–6.
34. Coiffier B, Neidhardt-Bérard EM, Tilly H, et al. Fludarabine alone compared to CHVP plus interferon in elderly patients with follicular lymphoma and adverse prognostic parameters: a GELA study. Groupe d'Etudes des Lymphomes de l'Adulte. *Annals in Oncology*. 1999;10:1191–7.
35. Bellosillo B, Villamor N, Colomer D, Pons G, Montserrat E, Gil J. In vitro evaluation of fludarabine in combination with cyclophosphamide and/or mitoxantrone in B-cell chronic lymphocytic leukemia. *Blood*. 1999;94:2836–43.
36. Alas S, Bonavida B, Emmanouilides C. Potentiation of fludarabine cytotoxicity on non-Hodgkin's lymphoma by pentoxifylline and rituximab. *Anticancer Research*. 2000;20(5A):2961–6.
37. Yamauchi T, Nowak BJ, Keating MJ, Plunkett W. DNA repair initiated in chronic lymphocytic leukemia lymphocytes by 4-hydroperoxycyclophosphamide is inhibited by fludarabine and clofarabine. *Clinical Cancer Research*. 2001;7:3580–9.
38. Di Gaetano N, Xiao Y, Erba E, et al. Synergism between fludarabine and rituximab revealed in a follicular lymphoma cell line resistant to the cytotoxic activity of either drug alone. *British Journal of Haematology*. 2001;114:800–9.
39. Priebe T, Platsoucas CD, Nelson JA. Adenosine receptors and modulation of natural killer cell activity by purine nucleosides. *Cancer Research*. 1990;50:4328–31.
40. Alas S, Emmanouilides C, Bonavida B. Inhibition of interleukin 10 by rituximab results in down-regulation of bcl-2 and sensitization of B-cell non-Hodgkin's lymphoma to apoptosis. *Clinical Cancer Research*. 2001;7:709–23.
41. DeVita VT Jr, Canellos GP, Chabner B, Schein P, Hubbard SP, Young RC. Advanced diffuse histiocytic lymphoma, a potentially curable disease. *Lancet*. 1975;1:248–50.
42. Fisher RI, Gaynor ER, Dahlborg S, et al. Comparison of a standard regimen (CHOP) with three intensive chemotherapy regimens for advanced non-Hodgkin's lymphoma. *New England Journal of Medicine*. 1993;328:1002–6.
43. Nickenig C, Dreyling M, Hoster E, et al. Combined cyclophosphamide, vincristine, doxorubicin, and prednisone (CHOP) improves response rates but not survival and has lower hematologic toxicity compared with combined mitoxantrone, chlorambucil, and prednisone (MCP) in follicular and mantle cell lymphomas: results of a prospective randomized trial of the German Low-Grade Lymphoma Study Group. *Cancer*. 2006;107:1014–22.
44. Hochster HS, Oken MM, Winter JN, et al. Phase I study of fludarabine plus cyclophosphamide in patients with previously untreated low-grade lymphoma: results and long-term follow-up—a report from the Eastern Cooperative Oncology Group. *Journal of Clinical Oncology*. 2000;18:987–94.
45. Flinn IW, Byrd JC, Morrison C, et al. Fludarabine and cyclophosphamide with filgrastim support in patients with previously untreated indolent lymphoid malignancies. *Blood*. 2000;96:71–5.
46. Hagemester F, Cabanillas F, Coleman M, Gregory SA, Zinzani PL. The role of mitoxantrone in the treatment of indolent lymphomas. *Oncologist*. 2005;10:150–9.
47. McLaughlin P, Hagemester FB, Romaguera JE, et al. Fludarabine, mitoxantrone, and dexamethasone: an effective new regimen for indolent lymphoma. *Journal of Clinical Oncology*. 1996;14:1262–8.
48. Zinzani PL, Bendandi M, Magagnoli M, Gherlinzoni F, Merla E, Tura S. Fludarabine-mitoxantrone combination-containing regimen in recurrent low-grade non-Hodgkin's lymphoma. *Annals in Oncology*. 1997;8:379–83.
49. Crawley CR, Foran JM, Gupta RK, et al. A phase II study to evaluate the combination of fludarabine, mitoxantrone and dexamethasone (FMD) in patients with follicular lymphoma. *Annals in Oncology*. 2000;11:861–5.
50. Velasquez WS, Lew D, Grogan TM, et al. Combination of fludarabine and mitoxantrone in untreated stages III and IV low-grade lymphoma: S9501. *Journal of Clinical Oncology*. 2003;21:1996–2003.
51. Zinzani PL, Magagnoli M, Moretti L, et al. Randomized trial of fludarabine versus fludarabine and idarubicin as frontline treatment in patients with indolent or mantle-cell lymphoma. *Journal of Clinical Oncology*. 2000;18:773–9.
52. Tsimberidou AM, McLaughlin P, Younes A, et al. Fludarabine, mitoxantrone, dexamethasone (FND) compared with an alternating triple therapy (ATT) regimen in patients with stage IV indolent lymphoma. *Blood*. 2002;100:4351–7.
53. Rohatgi N, LaRocca RV, Bard V, Sethuraman G, Foon KA. Phase II trial of sequential therapy with fludarabine followed by cyclophosphamide, mitoxantrone, vincristine, and prednisone for low-grade follicular lymphomas. *American Journal of Hematology*. 2002;70:181–5.
54. Foussard C, Colombat P, Maisonneuve H, et al. Long-term follow-up of a randomized trial of fludarabine-mitoxantrone, compared with cyclophosphamide, doxorubicin, vindesine, prednisone (CHVP), as first-line treatment of elderly patients with advanced, low-grade non-Hodgkin's lymphoma before the era of monoclonal antibodies. *Annals of Oncology*. 2005;16:466–72.
55. Cheson BD, Leonard JP. Monoclonal antibody therapy for B-cell non-Hodgkin's lymphoma. *New England Journal of Medicine*. 2008;359:613–26.
56. Colombat P, Salles G, Brousse N, et al. Rituximab (anti-CD20 monoclonal antibody) as single first-line therapy for patients with follicular lymphoma with a low tumor burden: clinical and molecular evaluation. *Blood*. 2001;97:101–6.
57. Czuczman MS, Grillo-López AJ, McLaughlin P, et al. Clearing of cells bearing the bcl-2 [t(14; 18)] translocation from blood and marrow of patients treated with rituximab alone or in combination with CHOP chemotherapy. *Annals of Oncology*. 2001;12:109–14.
58. Czuczman MS, Grillo-López AJ, White CA, et al. Treatment of patients with low-grade B-cell lymphoma with the combination of chimeric anti-CD20 monoclonal antibody and CHOP chemotherapy. *Journal of Clinical Oncology*. 1999;17:268–76.
59. Marcus R, Imrie K, Solal-Celigny P, et al. Phase III study of R-CVP compared with cyclophosphamide, vincristine, and prednisone alone in patients with previously untreated advanced follicular lymphoma. *Journal of Clinical Oncology*. 2008;26:4579–86.
60. McLaughlin P, Hagemester FB, Rodriguez MA, et al. Safety of fludarabine, mitoxantrone, and dexamethasone combined with rituximab in the treatment of stage IV indolent lymphoma. *Seminars in Oncology*. 2000;27(6 Suppl 12):37–41.
61. Czuczman MS, Koryzna A, Mohr A, et al. Rituximab in combination with fludarabine chemotherapy in low-grade or follicular lymphoma. *Journal of Clinical Oncology*. 2005;23:694–704.
62. Zinzani PL, Pulsoni A, Perrotti A, et al. Fludarabine plus mitoxantrone with and without rituximab versus CHOP with and without rituximab as front-line treatment for patients with follicular lymphoma. *Journal of Clinical Oncology*. 2004;22:2654–61.
63. Sacchi S, Pozzi S, Marcheselli R, et al. Rituximab in combination with fludarabine and cyclophosphamide in the treatment of patients with recurrent follicular lymphoma. *Cancer*. 2007;110:121–8.
64. Tam CS, O'Brien S, Wierda W, et al. Long-term results of the fludarabine, cyclophosphamide, and rituximab regimen as initial therapy of chronic lymphocytic leukemia. *Blood*. 2008;112:975–80.
65. Forstpointner R, Dreyling M, Repp R, et al. The addition of rituximab to a combination of fludarabine, cyclophosphamide, mitoxantrone (FCM) significantly increases the response rate and prolongs survival as compared with FCM alone in patients with relapsed and refractory follicular and mantle cell lymphomas: results of a prospective randomized study of the German Low-Grade Lymphoma Study Group. *Blood*. 2004;104:3064–71.

66. Forstpointner R, Unterhalt M, Dreyling M, et al. Maintenance therapy with rituximab leads to a significant prolongation of response duration after salvage therapy with a combination of rituximab, fludarabine, cyclophosphamide, and mitoxantrone (R-FCM) in patients with recurring and refractory follicular and mantle cell lymphomas: Results of a prospective randomized study of the German Low Grade Lymphoma Study Group (GLSG). *Blood*. 2006;108:4003–8.
67. Leonard JP, Coleman M, Kostakoglu L, et al. Abbreviated chemotherapy with fludarabine followed by tositumomab and iodine I¹³¹ tositumomab for untreated follicular lymphoma. *Journal of Clinical Oncology*. 2005;23:5696–704.
68. Bosch F, López-Guillermo A, Campo E, et al. Mantle cell lymphoma: presenting features, response to therapy, and prognostic factors. *Cancer*. 1998;82:567–75.
69. Decaudin D, Bosq J, Tertian G, et al. Phase II trial of fludarabine monophosphate in patients with mantle-cell lymphomas. *Journal of Clinical Oncology*. 1998;16:579–83.
70. Foran JM, Rohatiner AZ, Coiffier B, et al. Multicenter phase II study of fludarabine phosphate for patients with newly diagnosed lymphoplasmacytoid lymphoma, Waldenström's macroglobulinemia, and mantle-cell lymphoma. *Journal of Clinical Oncology*. 1999;17:546–53.
71. Kantarjian HM, Alexanian R, Koller CA, Kurzrock R, Keating MJ. Fludarabine therapy in macroglobulinemic lymphoma. *Blood*. 1990;75:1928–31.
72. Zinzani PL, Gherlinzoni F, Bendandi M, et al. Fludarabine treatment in resistant Waldenström's macroglobulinemia. *European Journal of Haematology*. 1995;54:120–3.
73. Leblond V, Ben-Othman T, Deconinck E, et al. Activity of fludarabine in previously treated Waldenström's macroglobulinemia: a report of 71 cases. Groupe Coopératif Macroglobulinémie. *Journal of Clinical Oncology*. 1998;16:2060–4.
74. Leblond V, Lévy V, Maloisel F, et al. Multicenter, randomized comparative trial of fludarabine and the combination of cyclophosphamide-doxorubicin-prednisone in 92 patients with Waldenström macroglobulinemia in first relapse or with primary refractory disease. *Blood*. 2001;98:2640–4.
75. Dhodapkar MV, Jacobson JL, Gertz MA, Crowley JJ, Barlogie B. Prognostic factors and response to fludarabine therapy in Waldenström's macroglobulinemia: an update of a U.S. intergroup trial (SW0G S9003). *Seminars in Oncology*. 2003;30:220–5.
76. Tam CS, Wolf MM, Westerman D, Januszewicz EH, Prince HM, Seymour JF. Fludarabine combination therapy is highly effective in first-line and salvage treatment of patients with Waldenström's macroglobulinemia. *Clinical Lymphoma and Myeloma*. 2005;6:136–9.
77. Treon SP, Branagan AR, Ioakimidis L, et al. Long term outcomes to fludarabine and rituximab in Waldenström macroglobulinemia. *Blood*. 2008 Nov 17. [Epub ahead of print]
78. Economopoulos T, Psyrris A, Fountzilas G, et al. Phase II study of low-grade non-Hodgkin lymphomas with fludarabine and mitoxantrone followed by rituximab consolidation: promising results in marginal zone lymphoma. *Leukemia Lymphoma*. 2008;49:68–74.
79. Foss FM, Ihde DC, Linnoila IR, et al. Phase II trial of fludarabine phosphate and interferon alfa-2a in advanced mycosis fungoides/Sézary syndrome. *Journal of Clinical Oncology*. 1994;12:2051–9.
80. Hast R, Jacobsson B, Petrescu A, Hjalmar V. Successful treatment with fludarabine in two cases of angioimmunoblastic lymphadenopathy with dysproteinemia. *Leukemia Lymphoma*. 1999;34:597–601.
81. Go RS, Gazelka H, Hogan JD, Wester SM. Subcutaneous panniculitis-like T-cell lymphoma: complete remission with fludarabine. *Annals of Hematology*. 2003;82:247–50.
82. Von Hoff DD, Dahlberg S, Hartstock RJ, Eyre HJ. Activity of fludarabine monophosphate in patients with advanced mycosis fungoides: a Southwest Oncology Group study. *Journal of the National Cancer Cancer*. 1990;82:1353–5.
83. Cheson BD. Infectious and immunosuppressive complications of purine analog therapy. *Journal of Clinical Oncology*. 1995;13:2431–48.
84. Eichhorst BF, Busch R, Schweighofer C, Wendtner CM, Emmerich B, Hallek M; German CLL Study Group (GCLLSG). Due to low infection rates no routine anti-infective prophylaxis is required in younger patients with chronic lymphocytic leukaemia during fludarabine-based first line therapy. *British Journal of Haematology*. 2007;136:63–72.
85. Picardi M, Pane F, Quintarelli C, et al. Hepatitis B virus reactivation after fludarabine-based regimens for indolent non-Hodgkin's lymphomas: high prevalence of acquired viral genomic mutations. *Haematologica*. 2003;88:1296–303.
86. Cheson BD, Vena DA, Foss FM, Sorensen JM. Neurotoxicity of purine analogs: a review. *Journal of Clinical Oncology*. 1994;12:2216–28.
87. Vidarsson B, Mosher DF, Salamat MS, Isaksson HJ, Onundarson PT. Progressive multifocal leukoencephalopathy after fludarabine therapy for low-grade lymphoproliferative disease. *American Journal of Hematology*. 2002;70:51–4.
88. Visani G, Lemoli RM, Tosi P, et al. Fludarabine-containing regimens severely impair peripheral blood stem cells mobilization and collection in acute myeloid leukaemia patients. *British Journal of Haematology*. 1999;105:775–9.
89. Laszlo D, Galieni P, Raspadori D, et al. Fludarabine containing-regimens may adversely affect peripheral blood stem cell collection in low-grade non Hodgkin lymphoma patients. *Leukemia Lymphoma*. 2000;37:157–61.
90. Michallet M, Thiébaud A, Dreger P, et al. Peripheral blood stem cell (PBSC) mobilization and transplantation after fludarabine therapy in chronic lymphocytic leukaemia (CLL): a report of the European Blood and Marrow Transplantation (EBMT) CLL subcommittee on behalf of the EBMT Chronic Leukaemias Working Party (CLWP). *British Journal of Haematology*. 2000;108:595–601.
91. Tam CS, Seymour JF, Prince HM, et al. Treatment-related myelodysplasia following fludarabine combination chemotherapy. *Haematologica*. 2006;91:1546–50.
92. McLaughlin P, Estey E, Glassman A, et al. Myelodysplasia and acute myeloid leukemia following therapy for indolent lymphoma with fludarabine, mitoxantrone, and dexamethasone (FND) plus rituximab and interferon alpha. *Blood*. 2005;105:4573–5.
93. Armitage JO, Carbone PP, Connors JM, Levine A, Bennett JM, Kroll S. Treatment-related myelodysplasia and acute leukemia in non-Hodgkin's lymphoma patients. *Journal of Clinical Oncology*. 2003;21:897–906.
94. Killick S, Mercieca J, Nandi A, Behrens J. Fludarabine-induced tumour lysis in low-grade non-Hodgkin's lymphoma. *Clinical and Laboratory Haematology*. 1997;19:79–80.
95. Weiss RB, Freiman J, Kweder SL, Diehl LF, Byrd JC. Hemolytic anemia after fludarabine therapy for chronic lymphocytic leukemia. *Journal of Clinical Oncology*. 1998;16:1885–9.
96. Borthakur G, O'Brien S, Wierda WG, et al. Immune anaemias in patients with chronic lymphocytic leukaemia treated with fludarabine, cyclophosphamide and rituximab—incidence and predictors. *British Journal of Haematology*. 2007;136:800–5.
97. Maung ZT, Wood AC, Jackson GH, Turner GE, Appleton AL, Hamilton PJ. Transfusion-associated graft-versus-host disease in fludarabine-treated B-chronic lymphocytic leukaemia. *British Journal of Haematology*. 1994;88:649–52.
98. Foran JM, Oscier D, Orchard J, et al. Pharmacokinetic study of single doses of oral fludarabine phosphate in patients with "low-grade" non-Hodgkin's lymphoma and B-cell chronic lymphocytic leukemia. *Journal of Clinical Oncology*. 1999;17:1574–9.
99. Oscier D, Orchard JA, Culligan D, et al. The bioavailability of oral fludarabine phosphate is unaffected by food. *Hematology Journal*. 2001;2:316–21.
100. Tobinai K, Watanabe T, Ogura M, et al. Phase II study of oral fludarabine phosphate in relapsed indolent B-Cell non-Hodgkin's lymphoma. *Journal of Clinical Oncology*. 2006;24:174–80.
101. Stefoni V, Alinari L, Musuraca G, et al. Efficacy and safety of oral fludarabine/cyclophosphamide regimen in previously treated indolent lymphomas. *Leukemia Lymphoma*. 2005;46:1839–41.

102. Boogaerts MA, Van Hoof A, Catovsky D, et al. Activity of oral fludarabine phosphate in previously treated chronic lymphocytic leukemia. *Journal of Clinical Oncology*. 2001;19:4252–8.
103. Fabbri A, Lenoci M, Gozzetti A, et al. Low-dose oral fludarabine plus cyclophosphamide as first-line treatment in elderly patients with indolent non-Hodgkin lymphoma. *British Journal of Haematology*. 2007;139:90–3.
104. Zinzani PL, Tani M, Pulsoni A, et al. Fludarabine and mitoxantrone followed by yttrium-90 ibritumomab tiuxetan in previously untreated patients with follicular non-Hodgkin lymphoma trial: a phase II non-randomised trial (FLUMIZ). *Lancet Oncology*. 2008; 9:352–8.
105. Zelenetz AD, Advani RH, Byrd JC, et al. National Comprehensive Cancer Network. Non-Hodgkin's lymphomas. *Journal of the National Comprehensive Cancer Network*. 2008;6:356–421. Available at: <http://www.nccn.org>. Accessed January 2009.
106. van Oers MH, Klasa R, Marcus RE, et al. Rituximab maintenance improves clinical outcome of relapsed/resistant follicular non-Hodgkin lymphoma in patients both with and without rituximab during induction: results of a prospective randomized phase 3 intergroup trial. *Blood*. 2006;108:3295–301.
107. Friedberg JW, Huang J, Dillon H, et al. National LymphoCare Study Investigators. Initial therapeutic strategy in follicular lymphoma (FL): An analysis from the National LymphoCare Study (NLCS). *Journal of Clinical Oncology*. 2006;24 (18 Suppl):7527.
108. Czuczman MS. Controversies in follicular lymphoma: “who, what, when, where, and why?” (Not necessarily in that order!). *Hematology*. 2006:303–10.
109. Yuen AR, Kamel OW, Halpern J, Horning SJ. Long-term survival after histologic transformation of low-grade follicular lymphoma. *Journal of Clinical Oncology*. 1995;13:1726–33.
110. Rodriguez J, McLaughlin P, Fayad L, et al. Follicular large cell lymphoma: long-term follow-up of 62 patients treated between 1973–1981. *Annals of Oncology*. 2000;11:1551–6.
111. Falkson CI. Fludarabine: a phase II trial in patients with previously treated low-grade lymphoma. *American Journal of Clinical Oncology*. 1996;19:268–70.
112. Fenchel K, Bergmann L, Wijermans P, et al. Clinical experience with fludarabine and its immunosuppressive effects in pretreated chronic lymphocytic leukemias and low-grade lymphomas. *Leukemia Lymphoma*. 1995;18:485–92.
113. Gillis S, Dann EJ, Cass Y, Rochlemer RR, Polliack A. Activity of fludarabine in refractory chronic lymphocytic leukemia and low grade non-Hodgkin's lymphoma—the Jerusalem experience. *Leukemia Lymphoma*. 1994;15:173–5.
114. O'Brien ME, Matutes E, Cunningham D, et al. Fludarabine in lymphoproliferative disorders: the Royal Marsden Hospital experience. *Leukemia Lymphoma*. 1994;14 (Suppl 2):17–23.
115. Bordonaro R, Petralia G, Restuccia N, et al. Fludarabine, mitoxantrone and dexamethasone as front-line therapy in elderly patients affected by newly-diagnosed, low-grade non-Hodgkin's lymphomas with unfavorable prognostic factors: results of a phase II study. *Leukemia Lymphoma*. 2004;45:93–100.
116. Cohen BJ, Moskowitz C, Straus D, Noy A, Hedrick E, Zelenetz A. Cyclophosphamide/fludarabine (CF) is active in the treatment of mantle cell lymphoma. *Leukemia Lymphoma*. 2001;42:1015–22.
117. Rohatgi N, LaRocca RV, Bard V, Sethuraman G, Foon KA. Phase II trial of sequential therapy with fludarabine followed by cyclophosphamide, mitoxantrone, vincristine, and prednisone for low-grade follicular lymphomas. *American Journal of Hematology*. 2002;70:181–5.
118. Cockcroft DW, Gault MH. Prediction of creatinine clearance from serum creatinine. *Nephron*. 1976; 16:31–41.

Appendix

Clinical trials using fludarabine currently conducted at the National Cancer Institute (available at: www.cancer.gov).

Phase III randomized study of fludarabine phosphate and cyclophosphamide with versus without rituximab in patients with previously untreated mantle cell lymphoma.

Rituximab after induction chemotherapy (R-CHOP vs. R-FC) followed by interferon maintenance versus rituximab maintenance in mantle cell lymphoma.

Phase III randomized study of first-line therapy comprising chlorambucil versus fludarabine phosphate in patients with previously untreated Waldenström macroglobulinemia, splenic lymphoma with villous lymphocytes, or non-IgM lymphoplasmacytic lymphoma.

Multicentric study, three randomized arms (R-CVP vs. R-CHOP vs. R-FM) for patients with stage II-IV follicular lymphoma [in Italy].

Phase I–II study of rituximab, fludarabine, and cyclophosphamide followed by radioimmunotherapy comprising rituximab and yttrium-90 ibritumomab tiuxetan in patients with relapsed indolent, mantle cell, or transformed CD20-positive B-cell non-Hodgkin's lymphoma.

Phase I–II study of haploidentical allogeneic aldesleukin-activated natural killer cells and aldesleukin following chemoimmunotherapy comprising rituximab, fludarabine, and cyclophosphamide in patients with CD20-positive relapsed non-Hodgkin lymphoma or chronic lymphocytic leukemia.

Rituximab, fludarabine, mitoxantrone, dexamethasone (R-FND) plus zevalin for high-risk (untreated) follicular lymphoma.

Bortezomib (Velcade) with standard chemotherapy (FMD-R) for relapsed or refractory follicular lymphoma.