

REVIEW

**OPEN ACCESS**  
Full open access to this and  
thousands of other papers at  
<http://www.la-press.com>.

## Montelukast: Pharmacology, Safety, Tolerability and Efficacy

Michael S. Benninger and Heather Waters

Head and Neck Institute, The Cleveland Clinic, Cleveland, OH, USA. Email: [benninm@ccf.org](mailto:benninm@ccf.org)

---

**Abstract:** Allergic rhinitis and asthma are common disorders effecting large percentages of the population of Western countries. There are multiple treatment options available for allergic rhinitis and asthma and stepwise approaches to therapy have been recommended. Montelukast is a cysteinyl leukotriene receptor antagonist that has been found to be effective both in the treatment of allergic rhinitis and asthma. This paper will describe the pharmacology, safety, efficacy and tolerability of montelukast. It will examine the comparative efficacy of montelukast to other medications for the treatment of allergic rhinitis and asthma, as well as discuss the recent studies of combination therapy.

**Keywords:** allergic rhinitis, asthma, montelukast

---

*Clinical Medicine: Therapeutics* 2009:1 1253–1261

This article is available from <http://www.la-press.com>.

© Libertas Academica Ltd.

This is an open access article distributed under the terms of the Creative Commons Attribution License (<http://www.creativecommons.org/licenses/by/2.0>) which permits unrestricted use, distribution and reproduction provided the original work is properly cited.

The authors grant exclusive rights to all commercial reproduction and distribution to Libertas Academica. Commercial reproduction and distribution rights are reserved by Libertas Academica. No unauthorised commercial use permitted without express consent of Libertas Academica. Contact [tom.hill@la-press.com](mailto:tom.hill@la-press.com) for further information.



## Introduction

Allergic rhinitis (AR) is a common disorder that affects up to 50–60 million persons in the U.S.,<sup>1</sup> accounting for 10%–40% of the population.<sup>2</sup> There is also strong evidence that the incidence may be increasing.<sup>3</sup> Given this high prevalence and the significant impact on quality of life, AR has been classified as a major chronic respiratory disease<sup>4</sup> and is now the 2nd leading cause of chronic disease in the United States.<sup>1</sup>


The characteristic symptoms of allergic rhinitis are nasal congestion, sneezing, anterior and posterior nasal drainage, nasal itching and itchy/red/watery eyes. Patients may also complain of throat clearing, facial discomfort or headache, postnasal drip, or itchy throat and/or palate.<sup>1,4–6</sup> Patients may also experience snoring, sleep disturbance, or fatigue and there is evidence that many patients may have decreased sense of sexual function.<sup>7</sup> The treatment of AR can lead to improvements in the specific allergic symptoms, enhancing overall quality of life and potentially reducing the incidence and/or severity of a number of comorbid disorders including asthma, rhinosinusitis, eustachian tube dysfunction and otitis media, allergic conjunctivitis, sleep disorders<sup>4,6,8</sup> and may improve sexual function.<sup>7</sup>

Asthma is also a very common disorder with 3.5% of the population in the United States having an asthmatic episode annually based on information from the US Department of Health and Human Services

and the Center for Disease Control.<sup>9</sup> The annual incidence of an asthmatic episode is increased in lower socio-economic groups.<sup>9</sup> Overall prevalence has been reported at approximately 15 million people or 1 person in 20 in the U.S.<sup>10</sup> These surveys also suggest that asthma is poorly controlled in the population as a whole with 9% of asthmatics requiring a hospitalization and 23% of asthmatic adults and 32% of asthmatic children requiring an emergency room visit.<sup>10</sup>

Evidence for the association between rhinitis and asthma has frequently been reported among epidemiologic studies. It has been stated that asthma may affect up to 40% of patients with allergic rhinitis, this percentage is significantly higher than that of the general population.<sup>11,12</sup> The prevalence rates of rhinitis among asthmatic patients can vary widely and can be as high as 80% depending on the age of onset. Rhinitis tends to precede the onset of asthma in patients with upper and lower airway disease and tends to be associated with bronchial hyperresponsiveness.<sup>11,13</sup> Findings suggest that rhinitis may be the initial manifestation of an allergic airway disease that leads to the onset of asthma.<sup>11,13,14</sup>

There have been a number of recommendations for a stepwise approach for the management of allergic rhinitis. Two highly regarded and well substantiated recommendations are the Practice Parameter recommendations<sup>6</sup> (Fig. 1) and the ARIA (Allergic Rhinitis and its Impact on Asthma)



**Figure 1.** Step-care therapy for allergic rhinitis: US 2008 Updated Practice Parameter recommendations (adapted from the Rhinitis Action Plan).<sup>6</sup>



guidelines.<sup>4</sup> In both of these recommendations there is a stepwise approach depending on the seasonality, duration and/or severity of disease. These also both include multiple different pharmacologic agents including oral antihistamines and decongestants, intranasal decongestants, intranasal corticosteroids, cromolyn sodium, intranasal antihistamines, and ipratropium bromide. One class of medication that has been found to be effective either by itself or in combination for the management of AR is the leukotriene modifiers. In the United States the leukotriene receptor antagonist (LTRA) approved for the treatment of AR is montelukast. Montelukast is also one of the approved LTRAs for the treatment of asthma. This report will describe the pharmacology and clinical trial data in relationship to the use of montelukast in AR and asthma. The literature was reviewed following a MEDLINE search for “montelukast, leukotriene receptor antagonists, cysteinyl leukotrienes, asthma and allergic rhinitis”. Review articles that describe AR and asthma were also evaluated. Articles were selected that describe key parameters of a review article on montelukast. Although a thorough search was accomplished, a systematic review of the literature was not performed.

### Cysteinyl Leukotrienes

Leukotriene LTC<sub>4</sub>, LTD<sub>4</sub>, LTE<sub>4</sub>, collectively termed cysteinyl leukotrienes (CysLTs) are peptide conjugated lipids that are the products of activated eosinophils, basophils, mast cells and macrophages.<sup>15</sup> CysLTs are arachidonic acid derivatives that are synthesized and released by immunocytes in the respiratory mucosa in response to the presence of an allergen.<sup>16–18</sup> They are appropriately named as the leukotrienes are produced by leukocytes and their chemical structure contains three double bonds, a triene.<sup>19</sup> The cysteinyl leukotrienes (C<sub>4</sub>, D<sub>4</sub>, E<sub>4</sub>) are generated from the arachidonic pathway via 5-lipoxygenase to intermediate leukotriene A<sub>4</sub>, then to leukotriene C<sub>4</sub> (by leukotriene c synthase), leukotriene D<sub>4</sub> (by gamma-glutamyl-transpeptidase) and leukotriene E<sub>4</sub> (by dipeptidase).<sup>20</sup> The cysteinyl leukotrienes are highly potent mediators of inflammation that lead to several reactions including contraction of bronchiolar smooth muscle, chemoattraction of eosinophils with stimulation of the release of other

mediators of inflammation, increased mucous production, and mucosal swelling brought about through increased vascular permeability and blood flow to the mucosa.<sup>16</sup> Increased CysLTs have been demonstrated in nasal lavage fluid in patients with allergic rhinitis after antigen provocation<sup>21,22</sup> and BAL washings in patients with asthma.<sup>23</sup> CysLTs exert their actions through activation of two G protein coupled receptors, termed CysLT subtype 1 receptor (CysLT1) and CysLT subtype 2 (CysLT2) receptor. CysLT1 is most studied and is the target for drugs such as montelukast.<sup>24</sup> Montelukast is a potent and highly selective antagonist of type 1 CysLT receptors, with an affinity two fold greater than the natural ligand.<sup>20,25</sup>

### Pharmacology

Montelukast is an orally active, highly selective LTRA that inhibits the CysLT1 receptor. Montelukast inhibits physiologic actions of LTD<sub>4</sub> at the CysLT1 receptor without any agonist activity. It is rapidly absorbed after administration reaching peak plasma concentration (C<sub>max</sub>) in 3 to 4 hours with a mean bioavailability of 64% following a 10 mg oral administration. For the 5 mg chewable tablet, the mean C<sub>max</sub> is achieved in 2 to 2.5 hours with a mean bioavailability of 73% fasting versus 63% with the standard meal. The 4 mg chewable tablet achieves C<sub>max</sub> in 2 hours, with fasting, in the 2 to 5 year old range.<sup>26</sup> More than 99% is bound to plasma proteins with minimal distribution across the blood-brain barrier. Metabolism occurs via liver P450 (CYP) 3A4 and 2C8 microsomes, with potent inhibition of P450 2C8. Excretion occurs almost exclusively in bile with a half-life from 2.7 to 5.5 hours in healthy adults. The pharmacokinetic profile is similar in females and males, young and elderly.<sup>26</sup>

In patients with mild to moderate hepatic insufficiency, no dosage adjustment is required but data is lacking regarding severe hepatic impairment. Montelukast and its metabolites are almost exclusively excreted in bile and not urine, and it therefore has not been evaluated in patients with renal insufficiency. In drug interaction studies, the recommended clinical dose of montelukast did not have clinically important effects on the pharmacokinetics of the following drugs: theophylline, prednisone, prednisolone, oral contraceptives, digoxin and



warfarin. Phenobarbital, which induces hepatic metabolism, decreased the plasma concentration of montelukast approximately 40% following a single 10 mg dose of montelukast, yet no dosage adjustment for montelukast is recommended in conjunction with phenobarbital usage.<sup>26</sup>

## Safety and Tolerability

Montelukast is well tolerated with a safety profile that is similar in adult and pediatric populations. Studies looking at safety and adverse effects have demonstrated no clinical or laboratory difference in adverse experiences versus placebo.<sup>16,27</sup> Adverse effects have been described as mild and most often include headache, gastrointestinal disturbances, fatigue, pharyngitis, upper respiratory tract infection and rash.<sup>16,19,27</sup> These events occurred at a frequency greater than or equal to 1% above placebo.<sup>26</sup> In a large phase IV study, 14 of 6158 patients (0.23%) reported 21 drug related adverse events consistent with the events noted above.<sup>28</sup>

Isolated reports of Churg-Strauss syndrome (CSS), a rare systemic vasculitis associated with asthma, have been described in asthma patients treated with montelukast.<sup>29,30</sup> CSS is a rare necrotizing vasculitis with an annual prevalence of approximately 60 per million in asthma patients and 2–7 per million in the general population.<sup>16,31</sup> Its precise etiology is not known, yet its clinical onset is most often preceded by discontinuation or tapering of corticosteroid therapy<sup>26,32</sup> and consists of eosinophilia, pulmonary infiltrates, cardiomyopathy and other signs of vasculitis.<sup>29</sup> A likely explanation is that inhaled or oral corticosteroid treatment in these patients may mask the underlying vasculitis that develops as the glucocorticosteroid doses are reduced in patients with severe asthma being treated with leukotriene receptor antagonists (LTRAs). Physicians should be aware of signs and symptoms of CSS in patients with moderate to severe asthma with tapering doses of corticosteroids.<sup>29</sup>

Although they are well tolerated, particular attention should be given to the consideration of periodically monitoring liver function tests during treatment with LTRA. There have been reported cases of liver dysfunction going from mild to severe in patients treated with these medications. Most hepatobiliary events occurred in combination with

other confounding factors, such as other medications or montelukast administration to patients with underlying potential for liver disease, such as alcohol use or hepatitis.<sup>26</sup> Elevations in alanine aminotransferase/glutamic oxaloacetic transaminase (ALT/GOT) and aspartate aminotransferase/glutamic pyruvic transaminase (AST/GPT) with montelukast occurred at a frequency that was generally comparable with placebo.<sup>26</sup> In patients treated with 10 mg as a single dose or duration up to 2 years, 2.1% had an increase in ALT compared to 2% of placebo. AST elevations were reported as adverse events in 1.6% and 1.2% of montelukast and placebo patients, respectively.<sup>26,29</sup> Consideration has been given to the fact that some patients who have demonstrated toxicity to LTRAs, possibly due to the toxic metabolite acyl-glucuronide, may have inherited a defect in the detoxification enzymes,<sup>29,33</sup> and therefore a genetic basis for drug-induced liver disease may exist.

Most recently, the FDA has published reports of agitation, aggression, anxiousness, dream abnormalities, hallucinations, depression, insomnia, irritability, restlessness, suicide, suicidal ideation, and tremor associated with the use of montelukast and other LTRAs based upon post marketing reports published by the drug manufacturer.<sup>34</sup> Wallerstedt et al. analyzed all reports of psychiatric disorders during treatment with montelukast in children occurring in Sweden. A total of 48 reports were found and included nightmares, anxiety, aggressiveness, sleep disorders, insomnia, irritability, hallucination, hyperactivity and personality disorder. Forty eight percent of these patients were 3 years of age or younger and in those where time to exposure was reported, 80% of these occurred within 1 week of beginning treatment.<sup>35</sup> Further studies have reported insufficient data to prove a link between montelukast and suicidality.<sup>36</sup> It has been reported that the rates of depression, anxiety, and sleep disturbance which are risk factors for suicide are greater in patients with allergic rhinitis compared to the general population and that patients with a history of allergy may have an increased rate of suicide.<sup>37</sup> Sleep disturbances have additionally been found to be more common in asthmatics and are associated with asthma control and quality of life.<sup>38</sup>



## Leukotriene Inhibiting Drug

Montelukast (Singulair) is one of the most commonly prescribed LTRAs available and the only LTRA approved for the treatment of both AR and asthma. This LTRA demonstrates rapid oral administration (three hours to peak plasma concentrations), near maximal plasma protein binding (99%) and excretion principally in the bile. The terminal half life of montelukast is 5 hours and is considered a pregnancy category B medication.<sup>26,39</sup>

Montelukast is indicated for adults and for children two years and above. It is available in tablet, chewable table and cherry flavored tablet form. Ten mg is recommended for ages 15 and older, 5 mg chewable tablets for patients 6 to 14 years of age, 4 mg chewable tablets in patients 2 to 5 years of age, and 4 mg oral granule formulation in pediatrics for asthma in 12 to 23 months and allergic rhinitis in 6 to 23 months of age.<sup>26</sup> Dosing is daily. Adult dosing can be taken with food, pediatric dosing should be avoided 1 hour before and 2 hours after meals. Montelukast is contraindicated in patients with acute asthma or those with severe hepatic impairment.<sup>26</sup> As human model studies in pregnant or nursing women have not been performed, therefore montelukast should only be used in this population if clearly indicated.<sup>26</sup> There are no drug interactions with oral contraceptive pills, warfarin, digoxin or theophylline. Coadministration with CYP 3A4 hepatic enzyme inducers (phenytoin, phenobarbital, rifampin) may decrease montelukast levels.<sup>26</sup>

## Clinical Trials Comparing Montelukast to Placebo

There have been a number of trials that support the efficacy of montelukast in comparison to placebo. A prospective, randomized, double-blind placebo controlled trial has shown that montelukast was superior to placebo in the management of sneezing, rhinorhea, itching and congestion in spring seasonal AR patients.<sup>40</sup> A study by Chervinsky et al showed similar improvement of symptoms in patients treated with montelukast in comparison to placebo in an evaluation of the results of three prospective clinical trials in fall AR patients.<sup>41</sup> A study in children 7 to 14 years of age showed a reduction in nasal and eye symptom scores and decreased blood eosinophil counts in subjects treated with

montelukast in comparison to placebo.<sup>42</sup> A randomized placebo-controlled trial has also shown efficacy of montelukast in comparison to placebo in total nasal symptom scores and a quality of life questionnaire in children 2–6 years of age.<sup>43</sup> Studies have also shown the effectiveness of montelukast in comparison to placebo in both seasonal and perennial AR patients in both adults and children as part of trials that evaluated montelukast in combination with other medications.<sup>44,45</sup> This improvement included the control of daily symptoms of nasal congestion, itching, sneezing and rhinorhea as well as other endpoints such as eye symptoms, global evaluations, nighttime symptom scores and quality of life surveys.<sup>44,45</sup>

The clinical efficacy of montelukast in adults and children with persistent asthma, including patients sensitive to aspirin, has been evaluated in several randomized, double-blind, placebo controlled trials. A study by Reiss et al demonstrated that montelukast improved airway obstruction ( $FEV_{1,}$  morning and evening peak expiratory flow rates) and patient reported end points (daytime asthma symptoms, as needed beta-agonist use, nocturnal awakenings) when compared with placebo in asthmatic patients greater than 15 years of age.<sup>46</sup> In over 12 weeks of treatment of patients aged 2 to 5 years, montelukast administered as a 4-mg chewable tablet produced significant improvements compared with placebo in multiple parameters of asthma control including: daytime asthma symptoms (cough, wheeze, trouble breathing, and activity limitation); overnight asthma symptoms (cough); the percentage of days with asthma symptoms; the percentage of days without asthma; the need for beta-agonist or oral corticosteroids; physician global evaluations; and peripheral blood eosinophils. The clinical benefit of montelukast was evident within 1 day of starting therapy.<sup>47</sup> Direct bronchoscopic exam after 6 weeks of montelukast therapy in adults demonstrated significantly reduced numbers of activated eosinophils and mast cells compared to placebo upon examination of bronchial mucosa.<sup>48</sup> Pizzichini et al have reported decreased sputum and blood eosinophils after 4 weeks of montelukast therapy compared to placebo.<sup>49</sup> In one study, montelukast therapy offered significantly greater protection against exercise-induced bronchoconstriction than placebo therapy as evidenced by the improvement in the area under the  $FEV_{1,}$  curve, the significant





improvement in the maximal decrease in FEV<sub>1</sub> after exercise and the time from the maximal decrease in FEV<sub>1</sub> to the return of lung function to within 5 percent of pre-exercise. After 12 weeks of treatment, patients in the montelukast group were more likely to rate their asthma control as better and less likely to require rescue therapy with a beta-agonist during or after exercise challenge.<sup>50</sup>

## Monteleukast in Comparison to other Medications

A number of studies have shown that intranasal corticosteroids are superior to montelukast in the management of symptoms of AR.<sup>51-54</sup> A double-blind randomized controlled trial showed that fluticasone propionate (FP) was superior to montelukast in the control of both daytime and nighttime symptoms as well as reduction in eosinophil counts in patients with seasonal AR.<sup>51</sup> In patients with asthma who were being treated with inhaled fluticasone propionate/salmeterol, the addition of FP provided superior rhinitis control to the addition of montelukast.<sup>52</sup> Two additional randomized, double-blind placebo controlled trials have also shown greater improvement in daytime and nighttime nasal symptom scores and individual nasal symptom scores.<sup>53,54</sup> A comparison of montelukast to pseudoephedrine showed comparable control of nasal symptom scores and quality of life scores and increasing peak nasal inspiratory flow rates in the treatment of AR.<sup>55</sup>

## Combination Therapy

There have been a number of studies that have compared montelukast in combination with non-sedating antihistamines (NSA) in comparison to montelukast alone, NSA alone or intranasal corticosteroids. In one study, a combination of montelukast and loratadine has been found to be superior to either montelukast or loratadine alone in seasonal allergic rhinitis<sup>45</sup> while in another study the combination of montelukast/loratadine was not significantly different from loratadine alone.<sup>44</sup> In patients with persistent AR, the combination of montelukast and either desloratadine or levocetirizine were more effective than monotherapy with either of these three medications.<sup>56</sup> In children 2-6 years old, montelukast was equivalent to cetirizine in the management of nasal symptoms of perennial AR

except for nasal itching which was better controlled with cetirizine.<sup>43</sup> Montelukast in combination with cetirizine showed better control of the clinical nasal symptoms and reduced eosinophil counts better than either montelukast or cetirizine alone in seasonal AR patients.<sup>57</sup> When given 6 weeks before the start of the pollen season this combination delayed the onset of allergic seasonal symptoms in comparison to monotherapy.<sup>57</sup> Once daily treatment of fexofenadine (120 mg) was found to be equivalent to montelukast (10 mg) plus loratadine (10 mg) on peak nasal inspiratory flow rates and symptoms in seasonal AR.<sup>58</sup> A study comparing fexofenadine-pseudoephedrine was equivalent to a combination loratadine-montelukast in control of symptoms quality of life questionnaire and peak inspiratory flow rates.<sup>59</sup> Combined montelukast and cetirizine showed a significant change from baseline in nasal symptoms and nasal flow rates.<sup>60</sup> Finally, a combination of montelukast with loratadine has been found to be superior to loratadine alone in the treatment of delayed pressure urticaria.<sup>61</sup>

Studies have been performed to evaluate the efficacy of an intranasal corticosteroid and combined montelukast with a NSA. In patients with both AR and asthma, both budesonide nasal spray and combined montelukast/cetirizine improved nasal symptoms and nasal airflow rates although had less substantial impact on laboratory measurements of nasal function such as rhinomanometry.<sup>62</sup> In a similar study comparing mometasone furoate nasal spray and combined montelukast/cetirizine, there was no difference between the groups for nasal symptoms, nasal flow and nasal laboratory tests.<sup>60</sup> When fluticasone propionate nasal spray was evaluated in relationship to a combination of loratadine/montelukast, median total nasal symptom scores were not statistically significantly different but scores in a quality of life questionnaire were better and the eosinophil count and eosinophilic cationic protein counts were lower in the fluticasone group.<sup>63</sup>

Combination therapy has also been assessed in patients with asthma. In a study comparing montelukast to budesonide alone (in 400 or 800 µg bid) and to combination montelukast/budesonide (400 µg bid), bronchial reactivity was improved in all of the budesonide groups in comparison to the montelukast



group while there were no differences between the budesonide groups.<sup>64</sup> Asthma maintenance and control may be improved with the addition of montelukast. One study demonstrated that the addition of montelukast to inhaled corticosteroid steroid (ICS) therapy alone or in combination with a long acting beta agonist in asthma patients resulted in a significant decrease in asthma attacks, emergency room visits, hospitalizations and oral corticosteroid use.<sup>65</sup> Several studies have shown that the addition of montelukast to ICS improves control of mild to moderate asthma compared to ICS therapy alone. Though salmeterol was clinically as effective, or even more effective than the addition of montelukast in these studies, montelukast may have a better long term safety profile.<sup>66,67</sup>

There are at least 2 studies that have shown that an intranasal glucocorticoid spray is more effective than a combination of montelukast with an NSA.<sup>51,68</sup> In a study of fluticasone propionate in comparison to montelukast/cetirizine, the fluticasone patients experienced greater improvement in nasal congestion and total nasal symptom scores.<sup>68</sup> The addition of montelukast or cetirizine to fluticasone propionate only improved nasal itching scores in comparison to fluticasone alone.<sup>68</sup> Similarly, fluticasone propionate and combination of montelukast/loratadine both improved daytime symptom scores in comparison to montelukast alone.<sup>51</sup> Fluticasone was, however, superior to either montelukast alone or the combination of montelukast/loratadine in the control of the nighttime symptom scores and the eosinophil count only was decreased in the fluticasone group.<sup>51</sup>

## Conclusion

Montelukast is an antileukotriene agent that impacts an important portion of the allergic inflammatory pathway. It has been shown to be an effective treatment for AR and in some extent for asthma. The effectiveness may be potentiated with the addition of a NSA in combination. The bulk of the evidence would suggest that montelukast is less effective than intranasal glucocorticoids for AR. When evaluating the results of studies comparing an INS to a combination of montelukast/NSA, the results are inconsistent, with some showing equivalence of treatment effects while others suggest that the INS may be more effective, particularly in nighttime symptoms.

## Disclosures

Dr Benninger

Consultant: GSK, Sanofe-Aventis, Schering, Alcon  
Speaker: Sanofe-Aventis, Alcon

Research Grants: GSK.

Dr Waters

No disclosures.

## References

1. Marple BF, Fornadley JA, Patel AA, et al. Keys to successful management of patients with allergic rhinitis: Focus on patient confidence, compliance, and satisfaction. *Otolaryngol Head Neck Surg.* 2007;136:S107–24.
2. Tripathi A, Patterson R. Impact of allergic rhinitis treatment on quality of life. *Pharmacoeconomics.* 2001;19:891–9.
3. US Department of Health and Human Services. Evidence Report/Technology Assessment AHRQ Publication No 02-E024 2002 May;54:1–198.
4. Bousquet J, Khaltaev N, Cruz AA, et al. Allergic Rhinitis and its Impact on Asthma (ARIA) 2008 Update. *Allergy.* 2008;63(Suppl 86):8–160.
5. Allergies in America. (Cited 2008 April 24) Available from <http://www.myallergiesinamericacom>.
6. Wallace DV, Dykewicz MS, Bernstein DI, et al. The diagnosis and management of rhinitis: An updated practice parameter. *J Allergy Clin Immunol.* 2008;122:S1–84.
7. Benninger MS, Benninger RM. The impact of allergic rhinitis on sexual activity, sleep and fatigue. *Allergy Asthma Proc.* 2009.
8. Storms WW. Rethinking our approach to allergic rhinitis management. *Ann Allergy Asthma Immunol.* 2002;88:30–5.
9. US Department of Health and Human Services. Health, United States. 2005:239.
10. Asthma in America. (Cited March 31, 2009) Available from <http://www.AsthmainAmericacom>.
11. Guerra S, Sherrill DL, Martinez FD, et al. Rhinitis as an independent risk factor for adult-onset asthma. *J Allergy Clin Immunol.* 2002;109:419–25.
12. Leynaert B, Neukirch F, Demoly P, et al. Epidemiologic evidence for asthma and rhinitis comorbidity. *J Allergy Clin Immunol.* 2000;106 (Suppl 5):S210–5.
13. Greisner WA, Settipane RJ, Settipane GA. Co-existence of asthma and allergic rhinitis: as 23 year follow-up study of college students. *Allergy Asthma Proc.* 1998;19:185–8.
14. Rusconi F, Galassi C, Corbo GM, et al. Risk factors for early, persistent and late-onset wheezing in young children. SIDRIA Collaborative group. *Am J Respir Crit Care Med.* 1999;160:1617–72.
15. Kanoaka Y, Boyce J. Cysteinyl leukotrienes and their receptors: cellular distribution and function in immune and inflammatory responses. *J Immunology.* 2004;173:1503–10.
16. Nayak A, Langdon R. Montelukast in the Treatment of Allergic Rhinitis. *Drugs.* 2007;67:887–90.
17. Creticos PS, Peters SP, Adkinson NF, et al. Peptide leukotriene release after antigen challenge in patients sensitive to ragweed. *N Engl J Med.* 1984;310:1626–30.
18. Volovitz B, Osur SL, Bernstein JM, et al. Leukotriene C4 release in upper respiratory mucosa during natural exposure to ragweed in ragweed-sensitive children. *J Allergy Clin Immunol.* 1988;82:414–8.
19. Dempsey OJ. Leukotriene receptor antagonist therapy. *Postgrad Med J.* 2000;76:767–73.
20. Lipworth BJ. Leukotriene-receptor antagonists. *Lancet.* 1999;353:57–62.
21. Fujita M, Nakagawa N, Yonetomi Y, et al. Cysteinyl leukotrienes induce nasal symptoms of allergic rhinitis via a receptor-mediated mechanism in guinea pigs. *Jpn J Pharmacol.* 1997;355–62.
22. Pipkorn U, Proud D, Lichtenstein LM, et al. Effect of short-term systemic glucocorticoid treatment on human nasal mediator release after antigen challenge. *J Clin Invest.* 1987;80:1957–61.



23. Busse W, Kraft M. Cysteinyl leukotrienes in allergic inflammation. *Chest*. 2005;127:1312–26.
24. Peters-Golden M, Gleason MM, Togias A. Cysteinyl leukotrienes: multi-functional mediators in allergic rhinitis. *Clin Exp All*. 2006;36:689–703.
25. Hay DW. Pharmacology of leukotriene receptor antagonists. More than inhibitors of bronchoconstriction. *Chest*. 1997;111(Suppl 2):35S–45S.
26. Merck & Co WS NJ. Singulair (montelukast sodium). Product information. (Cited 2009 April 20) Available from: [http://www.singulair.com/singulair/shared/documents/english/singulair\\_prescribing\\_infopdf](http://www.singulair.com/singulair/shared/documents/english/singulair_prescribing_infopdf) 2006.
27. Lagos JA, Marshall GD. Montelukast in the management of allergic rhinitis. *Ther Clin Risk Manag (Therapeutics and Clinical Risk Management)*. 2007;3:327–32.
28. Virchow JC, Bachert C RM. Efficacy and safety of montelukast in adults with asthma and allergic rhinitis. *Resp Med*. 2006;100:1952–9.
29. Riccioni G, Bucciarelli T, Mancini B, et al. Antileukotriene drugs: Clinical application, effectiveness and safety. *Cur Med Chem (Current Medical Chemistry)*. 2007;14:1966–77.
30. Wechsler ME, Finn D, Gunawardena D, et al. Churg-Strauss syndrome in patients receiving montelukast as treatment for asthma. *Chest*. 2000;117:708–13.
31. Loughlin JE, Cole JA, Rothman KJ, et al. Prevalence of serious eosinophilia and incidence of Churg-Strauss syndrome in a cohort of asthma patients. *Ann Allergy Asthma Immunol*. 2002;88:319–25.
32. Weller PF, Plaut M, Taggart V, et al. The relationship of asthma therapy and Churg-Strauss syndrome: NIH workup summary report. *J Allergy Clin Immunol*. 2001;108:175–83.
33. Actis GC, Morgando A, Laget M, et al. Zafirlukast-related hepatitis: report of a further case. *J Hepatol*. 2001;35:539–41.
34. FDA. (Cited 2009, July 15) Available from: <http://www.fda.gov/Drugs/DrugSafety/PostmarketDrugSafetyInformationforPatientsandProviders/DrugSafetyInformationforHealthcareProfessionals/ucm165489.htm>.
35. Wallerstedt SM, Brunlof G, Sunderstrom A, et al. Montelukast and psychiatric disorders in children. *Pharmacoepidemiol Drug Saf*. 2009 June 23: Pre-publication.
36. Manalai P, Woo JM, Postolache TT. Suicidality and montelukast. *Expert Opin Drug Saf*. 2009;8:273–82.
37. Postolache TT, Komarow H, Tonelli LH. Allergy: a risk factor for suicide? *Curr Treat Options Neurol*. 2008;10:363–76.
38. Mastrorade JG, Wise RA, Shade DM, et al. American Lung Association Asthma Clinical Research Centers. *J Asthma*. 2008;45:183–9.
39. AstraZeneca Pharmaceuticals W, Del. Accolate (zafirlukast) Product information. (Cited 2009, April 20) Available from: <http://www.astrazeneca-us.com/pi/accolatepdf> 2004.
40. Philip G, Malmstrom K, Hampel FC, et al. Montelukast for treating seasonal allergic rhinitis: a randomized, double-blind, placebo-controlled trial performed in the spring. *Clin Exp Allergy*. 2002;32:1020–8.
41. Chervinsky P, Philip G, Malice MP, et al. Montelukast for treating fall allergic rhinitis: effect of pollen exposure in 3 studies. *Ann Allergy Asthma Immunol*. 2004;92:367–73.
42. Razi C, Bakirtas RC, Harmanci K, et al. Effect of montelukast on symptoms and exhaled nitric oxide levels in 7- to 14-year-old children with seasonal allergic rhinitis. *Ann Allergy Asthma Immunol*. 2006;97:767–74.
43. Chen ST, Lu KH, Sun HL, et al. Randomized placebo-controlled trial comparing montelukast and cetirizine for treating perennial allergic rhinitis in children aged 2–6 years. *Pediatr Allergy Immunol*. 2006;17:4–53.
44. Nayak AS, Philip G, Lu S, et al. Efficacy and tolerability of montelukast alone or in combination with loratadine in seasonal allergic rhinitis: a multicenter, randomized, double-blind, placebo-controlled trial in the fall. *Ann Allergy Asthma Immunol*. 2002;88:592–600.
45. Meltzer EO, Malmstrom K, Lu S, et al. Concomitant montelukast and loratadine as treatment for seasonal allergic rhinitis: A randomized, placebo-controlled clinical trial. *J Allergy Clin Immunol*. 2000;105:917–22.
46. Reiss TF, Chervinsky P, Dockhorn RJ, et al. Montelukast, a once-daily leukotriene receptor antagonist, in the treatment of chronic asthma: a multicenter, randomized, double-blind trial. Montelukast Clinical Research Study Group. *Arch Intern Med*. 1998;158:1213–20.
47. Knorr B, Franchi LM, Bisgaard H, et al. Montelukast, a leukotriene receptor antagonist, for the treatment of persistent asthma in children aged 2 to 5 years. *Pediatrics*. 2001;108:E48.
48. Ramsay CF, Sullivan P, Gizycki M, et al. Montelukast and bronchial inflammation in asthma: a randomized double-blind placebo-controlled trial. *Respir Med*. 2009;103:995–1003.
49. Pizzichini E, Leff JA, Reiss TF, et al. Montelukast reduces airway eosinophilic inflammation in asthma: a randomized, controlled trial. *Eur Respir J*. 1999;14:12–8.
50. Leff JA, Busse WW, Pearlman D, et al. Montelukast, a leukotriene-receptor antagonist, for the treatment of mild asthma and exercise-induced bronchoconstriction. *N Engl J Med*. 1998;339:147–52.
51. Pullerits T, Praks L, Ristioja V, et al. Comparison of a nasal glucocorticoid, antileukotriene, and a combination of antileukotriene and antihistamine in the treatment of seasonal allergic rhinitis. *J Allergy Clin Immunol*. 2002;109:949–55.
52. Nathan RA, Yancy SW, Waitkus-Edwards K, et al. Fluticasone propionate nasal spray is superior to montelukast for allergic rhinitis while neither affects overall asthma control. *Chest*. 2005;128:1910–20.
53. Ratner PH, Howland WC, Arustu R, et al. Fluticasone propionate Aqueous nasal spray provided significantly greater improvement in daytime and nighttime nasal symptoms of seasonal allergic rhinitis compared with montelukast. *Ann Allergy Asthma Immunol*. 2003;2003:466–8.
54. Martin BG, Andrews CP, van Bavel JH, et al. Comparison of fluticasone propionate aqueous nasal spray and oral montelukast for the treatment of seasonal allergic rhinitis symptoms. *Ann Allergy Asthma Immunol*. 2006;96:851–7.
55. Mucha SM, deTineo M, Naclario RM, et al. Comparison of montelukast and pseudoephedrine in the treatment of allergic rhinitis. *Arch Otolaryngol Head Neck Surg*. 2006;132:164–72.
56. Ciebiada M, Gorska-Ciebiada M, DuBuske LM, et al. Montelukast with desloratadine or levocetirizine for the treatment of persistent allergic rhinitis. *Ann Allergy Asthma Immunol*. 2006;97:664–71.
57. Kurowski M, Kuna P, Gorski P. Montelukast plus cetirizine in the prophylactic treatment of seasonal allergic rhinitis: influence on clinical symptoms and nasal allergy inflammation. *Allergy*. 2004;58:280–8.
58. Wilson AM, Orr LC, Coutie WJR, et al. A comparison of once daily fexofenadine versus the combination of montelukast plus loratadine on domiciliary nasal peak flow and symptoms in seasonal allergic rhinitis. *Clin Exp Allergy*. 2002;32:126–32.
59. Moinuddin R, deTineo M, Maleckar B, et al. Comparison of the combination of fexofenadine-pseudoephedrine and loratadine-montelukast in the treatment of seasonal allergic rhinitis. *Ann Allergy Asthma Immunol*. 2004;92:73–9.
60. Wilson AM, Orr LC, Sims EJ, et al. Effects of monotherapy with intra-nasal corticosteroid or combined oral antihistamine and leukotriene receptor antagonists in seasonal allergic rhinitis. *Clin Exp Allergy*. 2001;31:61–8.
61. Nettis E, Pannofino A, Corvello E, et al. Efficacy of montelukast in combination with loratadine, in the treatment of delayed pressure urticaria. *J Allergy Clin Immunol*. 2003;112:212–3.
62. Wilson AM, Sims EJ, Orr LC, et al. Effects of topical corticosteroid and combined mediator blockade on domiciliary and laboratory measurements of nasal function in seasonal allergic rhinitis. *Ann Allergy Asthma Immunol*. 2001;87:249–344.
63. Saengpanich S, deTineo M, Naclario RM, et al. Fluticasone nasal spray and the combination of loratadine and montelukast in seasonal allergic rhinitis. *Arch Otolaryngol Head Neck Surg*. 2003;129:557–62.
64. Riccioni G, Vecchia RD, Orazio ND, et al. Comparison of montelukast and budesonide on bronchial reactivity in subjects with mild-moderate persistent asthma. *Pulm Pharm Ther*. 2002;16:111–4.





65. Borderias L, Mincewicz G, Paggiaro PL, et al. Asthma control in patients with asthma and allergic rhinitis receiving add-on montelukast therapy for 12 months: a retrospective observation study. *Curr Med Res Opin.* 2007; 23:721–30.
66. Joos S, Miksch A, Szecsenyi J, et al. Montelukast as an add-on therapy to inhaled corticosteroids in the treatment of mild to moderate asthma: a systemic review. *Thorax.* 2008;63:453–62.
67. Ducharme FM, Lasserson TJ, Cates CJ. Long-acting beta2-agonists versus anti-leukotrienes as add-on therapy to inhaled corticosteroids for chronic asthma. *Cochrane Database Syst Rev.* 2006 Oct 18;(4):CD003137.
68. DiLorenza G, Pacort ML, Pellitieri ME, et al. Randomized placebo-controlled trial comparing fluticasone aqueous nasal spray in mono-therapy, fluticasone plus cetirizine, fluticasone plus montelukast and cetirizine plus montelukast for seasonal allergic rhinitis. *Clin Exp Allergy.* 2004;34:259–67.

**Publish with Libertas Academica and every scientist working in your field can read your article**

*“I would like to say that this is the most author-friendly editing process I have experienced in over 150 publications. Thank you most sincerely.”*

*“The communication between your staff and me has been terrific. Whenever progress is made with the manuscript, I receive notice. Quite honestly, I’ve never had such complete communication with a journal.”*

*“LA is different, and hopefully represents a kind of scientific publication machinery that removes the hurdles from free flow of scientific thought.”*

**Your paper will be:**

- Available to your entire community free of charge
- Fairly and quickly peer reviewed
- Yours! You retain copyright

**<http://www.la-press.com>**