

REVIEW

OPEN ACCESS

Full open access to this and thousands of other papers at <http://www.la-press.com>.

A Review of Sevelamer Hydrochloride in End-Stage Renal Disease Patients on Dialysis

Goce Spasovski

Department of Nephrology, University of Skopje, Skopje, R. Macedonia. Corresponding author email: gspas@sonet.com.mk

Abstract: It is known that in the presence of even subtle kidney dysfunction an intensive prevention of cardiovascular risk is required. Apart from the conventional factors which contribute to cardiovascular disease (CVD), there are also some specific conditions of the chronic kidney disease (CKD) population such as oxidative stress of uremia and dialysis (D). However, hyperphosphatemia, hypercalcemia, and elevated calcium-phosphorus product remain as major contributors to the development of vascular calcification (VC) in this population, as part of the systemic complication known as mineral and bone disorders (MBD) in CKD patients. Importantly, the retention of phosphate remains as main culprit in the pathogenesis of CKD—MBD. Over the years, various treatment options for phosphate removal and controlling mineral metabolism, bone health, VC and CVD have failed, mainly through an over-suppression of PTH, development of ABD and promotion of VC and mortality.

Although KDOQI and KDIGO published CKD—MBD guidelines has clearly stated where calcium-based phosphate binders should not be used in D patients (hypercalcemia and low PTH) and where non calcium-containing phosphate binders are preferred (patients with severe vascular and/or other soft tissue calcifications), the greatest controversy and disagreements within the nephrological community still exists upon the cost-effectiveness of non calcium binder (sevelamer) use. Indeed, despite the evidence and recognised trend towards both a decrease in VC and CVD associated with sevelamer use, it is still an ongoing matter of debate. The magnitude of this controversy is increased when the issue of advanced medical and/or budgetary evaluation related to the implementation of clinical guidelines for CKD—MBD treatment is considered. Despite advocated use of sevelamer across a range of common clinical scenarios in CKD, its widespread utilization is challenged as exceeding what would usually be considered good value for money. If so, it is questionable whether the recommendations and suggestions from the guidelines should be followed, and further, do we need guidelines and innovative drugs for treatment of hyperphosphatemia? While awaiting the answer, as clinicians we should proceed with a treatment to “do no harm”, trying to at least limit the calcium exposure of our dialysis patients.

Keywords: hyperphosphatemia, KDOQI guidelines, KDIGO guidelines, calcium based binders, non calcium based binders, sevelamer

Clinical Medicine Insights: Therapeutics 2011:3 51–55

doi: [10.4137/CMT.S5990](https://doi.org/10.4137/CMT.S5990)

This article is available from <http://www.la-press.com>.

© the author(s), publisher and licensee Libertas Academica Ltd.

This is an open access article. Unrestricted non-commercial use is permitted provided the original work is properly cited.

The authors grant exclusive rights to all commercial reproduction and distribution to Libertas Academica. Commercial reproduction and distribution rights are reserved by Libertas Academica. No unauthorised commercial use permitted without express consent of Libertas Academica. Contact tom.hill@la-press.com for further information.



Introduction

Because there is an undeniable link between kidney dysfunction and cardiovascular risk, the presence of even subtle kidney dysfunction should be considered as one of the conditions necessitating intensive prevention of such risk.¹ Apart from the conventional factors such as hypertension, dyslipidemia, and hyperhomocysteinemia which contribute to cardiovascular disease (CVD), there are also some specific conditions of the chronic kidney disease (CKD) population such as oxidative stress of uremia and hemodialysis.² However, hyperphosphatemia, hypercalcemia, and elevated calcium-phosphorus product remain as major contributors to the development of vascular calcification (VC) in this population.^{3,4} This issue should also be looked at from a perspective of the common clinically defined systemic complication known as mineral and bone disorders (MBD) in CKD patients.⁵ MBD consists of a combination of mineral, hormonal and bone abnormalities, as well as vascular and soft tissue calcifications paralleling progressive kidney dysfunction. Of note, the retention of phosphate is considered as the main culprit in the pathogenesis of MBD in patients with advanced CKD.⁶ In addition, the production of fibroblast growth factor 23 (FGF-23), a novel bone-derived phosphaturic hormone that inhibits both renal phosphate reabsorption and calcitriol production, should be regarded as an important player in CKD-MBD.⁷ Here, the decreased production of calcitriol is a negative signal for FGF23 production basically stimulating parathyroid hormone (PTH) secretion, which in turn increases relative phosphate excretion despite the reduction in glomerular filtration rate (GFR). On the other hand, the reduced GFR stimulates production of FGF-23 levels which activates the bone-kidney axis coordinating systemic phosphate homeostasis and bone mineralization to protect the body from hyperphosphatemia. Although FGF23 could not differentiate between bone biopsy diagnosed low- or high-turnover bone disease,⁸ the beneficial effects of lowering FGF23 levels are suggested by the correlation between FGF23, vascular calcification, CKD progression and mortality.^{9,10} In summary FGF23 has uncovered new regulatory pathways and system biology governing mineralization, vitamin D metabolism, parathyroid gland function, and renal phosphate handling. Thus FGF23 assessment will become important in diagnosing of hypo- and hyperphosphatemic

disorders, for which pharmacological regulation of FGF23 levels may provide novel treatments.¹¹

Over the years various treatment options for phosphate removal or reduction were adopted in routine clinical practice.¹² Unfortunately, some treatment modalities to control mineral metabolism, bone health, VC and CVD have failed.¹³ Thus, the great expectations for calcium based phosphate binders were dissolved in the last decade when enhanced risk for over-suppression of parathyroid hormone (PTH) and development of adynamic bone disease (ABD), especially when used in combination with vitamin D, promoting VC and mortality became apparent.¹⁴⁻¹⁶ Indeed, hyperphosphatemia has been considered as one of the most “expensive” complications in treatment of CKD population. The aim of this review is to provide better insight in the newer treatment of hyperphosphatemia based on the evidence and surrounding controversy for its cost-efficiency.

Guidelines for Treatment of Hypersphosphatemia

The National Kidney Foundation Kidney Disease Outcomes Quality Initiative (KDOQI) has published the first Bone Metabolism and Disease Treatment Guidelines in 2003, in line with the growing body of evidence linking various treatment strategies and related clinical outcomes, following their goal of improving the quality of care and outcomes of patients with kidney disease.¹⁷ In the guidelines 5.6 (evidence) and 5.7 (opinion based) it is clearly stated where calcium-based phosphate binders should not be used in dialysis patients (hypercalcemia and low PTH levels below 150 pg/mL on 2 consecutive measurements)^{18,19} and where noncalcium-containing phosphate binders are preferred (patients with severe vascular and/or other soft tissue calcifications).²⁰

Although the authors of these guidelines acknowledged that additional hard evidence is needed to complete future revisions to these guidelines, the low evidence ‘judgements’ presented in the guidelines was generally perceived as much as absolute truth by the medical community as it is in the high evidence ‘guidelines’.²¹ Considering the bias of the existing evidence especially in the field of newer therapeutics and related clinical outcomes KDIGO initiative has recently launched the new guidelines on CKD-MBD.²² In the presence of not conclusive evidence of non- vs.



calcium-based binders, KDIGO recommended and suggested in guidelines 4.1.5 (1B) and (2C) limited calcium intake in the form of phosphate binders as more beneficial than harmful until further research is available. Repeatedly the restriction was proposed in the presence of arterial calcification, ABD and/or persistently low serum PTH levels and hypercalcemia in addition to stopped therapy with vitamin D. So both KDOQI and recent KDIGO guidelines were not different with respect to the limited treatment with calcium based binders. In addition, KDIGO guidelines pointed to at least some evidence in humans showing beneficial effects of sevelamer-HCl compared with calcium-based binders mainly on progression of arterial calcification and partially on mortality risk reduction. It should also be noted that there was no consensus for this guideline production from all the KDIGO members, considering its potential impact as too large in the presence of the scarce evidence to support it.

Sevelamer Hydrochloride in Treatment of CKD Patients— What is the Evidence?

The management of hyperphosphatemia and other components of MBD may be evaluated through various surrogate clinical endpoints (vascular calcification and bone disease), or hard clinical outcomes (cardiovascular events). Although a number of newer therapeutics are already available: Renagel® and Renvela® (Genzyme Corporation, Cambridge, MA); Fosrenol® (Shire Pharmaceuticals, Hampshire, UK); Sensipar® and Mimpara® (Amgen, Inc., Thousand Oaks, CA) to facilitate the achievement of consistent control of multiple MBD parameters, the majority of beneficial outcome data were reported only for sevelamer use.¹² At the same time, the greatest controversy and disagreements within the nephrological community continue on the cost-effectiveness of sevelamer use, especially during the global economic and medicare crisis, in both developed and developing countries.

What is the real evidence?

Considering the limitations of data generated from existing meta-analysis,²³ the two systematic reviews comparing sevelamer to other therapies could not find convincing evidence that sevelamer improves clinically relevant outcomes in ESRD patients,²⁴ pronouncing it as economically unattractive treatment strategy.²⁵

Later data on mortality benefits with sevelamer treatment emerged from randomised clinical trials. Namely, Block et al demonstrated significant survival benefit (as a secondary endpoint) in incident dialysis patients receiving sevelamer vs. calcium-based binders in a relatively small trial of 127 subjects (11 vs. 23 deaths, respectively).²⁶ On the other hand, the expectations in the largest outcome study ever conducted in the prevalent dialysis population (Dialysis Clinical Outcomes Revisited—DCOR) were not confirmed for all-cause mortality in the overall population.²⁷ Nevertheless, in a specified subgroup analysis of older population (age >65 years) and in patients treated with sevelamer for more than 2 years, sevelamer treatment was associated with a lower all-cause mortality. A possible difference in the reports from these two studies may be related to the various study designs (age, diabetes, dialysis duration, type of the study population, incident vs. prevalent) and the shorter follow up in the DCOR trial. Of note, in a secondary analysis of the DCOR study, there was evidence that sevelamer treated patients were less frequently (11%) hospitalized and spent less time (12%) in hospital.²⁸ In addition, beneficial effects of sevelamer treatment with increased bone formation and improved trabecular architecture although without statistically significant changes in bone turnover or mineralization compared with calcium carbonate were recently reported in a single 1-year bone biopsy based study.²⁹

In spite of all the above mentioned evidence and a recognised trend towards both a decrease in cardiovascular mortality and all measures of coronary artery calcification, in a two recent meta-analyses the decrease in mortality associated with sevelamer was still questioned as controversial matter.^{30,31} In fact, the authors could not conclude whether the beneficial effects came from an associated decrease in cholesterol, a decrease in coronary artery calcification, other pleiotropic effects of sevelamer or contrarily, an increase in mortality associated with calcium-based phosphate binders (CBPB).³¹

Benefits of Treatment with Sevelamer Compared to Calcium Based Binders

There is a plenty of new evidence with significantly lower coronary artery calcification scores in prevalent dialysis patients treated with sevelamer as compared to CBPB.³² Most probably besides the reduced calcium loading, there is an additional effect of sevelamer



increasing the calcification inhibitor levels of fetuin-A.³³ Other pleiotropic effects which may play a key role in the vascular protective activities of sevelamer as shown in a recent meta-analysis are lowering of the C-reactive protein levels, and a higher alkaline phosphatase and intact parathyroid hormone levels found among sevelamer-treated patients.³⁴ Thus, although calcium-based binders and sevelamer are almost equally effective for treatment of hyperphosphatemia, it is obvious that both phosphate binders have different mechanisms of action. The main adverse effect of CBPBs is the calcium excess accumulated through the use of binders, supplements, and dialysate which can lead to hypercalcemia, contribute to VC, and potentially affect bone histology and mortality. In contrast, sevelamer treatment (once or three times daily) results only in mild gastrointestinal adverse effect.³⁵

Recently, a new combined phosphate binder calcium acetate/magnesium carbonate (CaMg) has been offered as a therapeutic option, non-inferior in comparison with sevelamer at controlling serum phosphorus levels.³⁶ However, this report was challenged assuming that one (phosphate) CKD-MBD variable was well controlled in the CaMg group vs. total of four in the sevelamer group (calcium, PTH, magnesium). Additionally, the control of LDL cholesterol and potassium levels which was superior under sevelamer-treatment should have been taken into account as an important factor in survival outcomes.³⁷

Despite all existing data, there is still uncertainty in the nephrological community that the most cost-effective way to treat hyperphosphatemia in patients with end-stage renal disease is to be determined.³⁸ The magnitude of controversy is increased when the issue of advanced medical and/or budgetary evaluation related to the implementation of clinical guidelines for CKD—MBD treatment is considered. On top of it, the growing number of patients requiring dialysis and especially the high cost of CKD-MBD treatment itself are pressing clinicians to pharmaco-economically justify the management of hyperphosphatemia with new drugs available on the market and related outcomes. Thus, despite of the advocated use of sevelamer across a range of common clinical scenarios in CKD²² and demonstrated evidence of reduced morbidity and mortality,^{27,28} its widespread utilization was challenged as exceeding what would usually be considered good value for the money.^{39,40}

In conclusion, the controversy with regard to the sevelamer treatment seems to be against compelling adoption of the K/DOQI recommendations today and those of KDIGO's in the future. The extrapolated conclusion which should be drawn here is about the assessment of the worthwhile medical values which should be implemented for the improved well being of our patients. Medical professionals need an answer to the question whether they should follow recommendations and suggestions from the guidelines or if economic constraints should take precedence? Finally, one could ask, do we need guidelines and innovative drugs for treatment of hyperphosphatemia, if there is no possibility that they can be implemented in everyday clinical practice? Unfortunately, while awaiting answers, the only clinical perspective for treatment is to “do no harm”, trying to at least limit the calcium exposure of dialysis patients.

Disclosure

This manuscript has been read and approved by the author. This paper is unique and is not under consideration by any other publication and has not been published elsewhere. Goce Spasovski declares having received research grants from Shire, Fresenius Medical Care, Amgen and Mitsubishi Pharma, and honoraria as a speaker from Hoffmann-La Roche, Genzyme and Novartis. The peer reviewers of this paper report no conflicts of interest. The author confirms that they have permission to reproduce any copyrighted material.

References

1. Vanholder R, Massy Z, Argiles A, Spasovski G, et al. Chronic kidney disease as cause of cardiovascular morbidity and mortality. *Nephrol Dial Transplant*. 2005;20(6):1048–56.
2. Locatelli F, Canaud B, Eckardt KU, Stenvinkel P, Wanner C, Zoccali C. Oxidative stress in end-stage renal disease: an emerging threat to patient outcome. *Nephrol Dial Transplant*. 2003;18:1272–80.
3. Block GA, Klassen PS, Lazarus JM, et al. Mineral metabolism, mortality, and morbidity in maintenance hemodialysis. *J Am Soc Nephrol*. 2004;15:2208–18.
4. Vanholder R, Abou-Deif O, Argiles A, et al. The role of EUTox in uremic toxin research. *Semin Dial*. 2009;22(4):323–8.
5. Moe S, Drueke T, Cunningham J, et al. Kidney Disease: Improving Global Outcomes (KDIGO). Definition, evaluation, and classification of renal osteodystrophy: A position statement from Kidney Disease: Improving Global Outcomes (KDIGO). *Kidney Int*. 2006;69:1945–53.
6. Spasovski G, Massy Z, Vanholder R. Phosphate metabolism in chronic kidney disease: from pathophysiology to clinical management. *Semin Dial*. 2009;22(4):357–62.
7. Liu S, Tang W, Zhou J, et al. Fibroblast growth factor 23 is a counter-regulatory phosphaturic hormone for vitamin D. *J Am Soc Nephrol*. 2006;17:1305–15.
8. Cancela AL, Oliveira RB, Gracioli FG, et al. Fibroblast growth factor 23 in hemodialysis patients: effects of phosphate binder, calcitriol and calcium concentration in the dialysate. *Nephron Clin Pract*. 2010;117(1):c74–82.



9. Jean G, Bresson E, Terrat JC, et al. Peripheral vascular calcification in long-haemodialysis patients: Associated factors and survival consequences. *Nephrol Dial Transplant*. 2008;24:948–55.
10. Gutierrez OM, Mannstadt M, Isakova T, et al. Fibroblast growth factor 23 and mortality among patients undergoing hemodialysis. *N Engl J Med*. 2008;359:584–92.
11. Liu S, Gupta A, Quarles SO. Emerging role of fibroblast growth factor 23 in a bone-kidney axis regulating systemic phosphate homeostasis and extracellular matrix mineralization. *Curr Opin Nephrol Hypertens*. 2007;16(4):329–35.
12. Spasovski G. New strategies in treatment of mineral and bone disorders and associated cardiovascular disease in patients with chronic kidney disease. *Recent Patents Cardiovasc Drug Discov*. 2008;3(3):222–8.
13. Spasovski GB. Bone health and vascular calcification relationships in chronic kidney disease. *Int Urol Nephrol*. 2007;39(4):1209–16.
14. Goodman WG, Goldin SJ, Kuizon BD. Coronary artery calcification in young adults with end-stage renal disease who are undergoing dialysis. *N Engl J Med*. 2000;342:1478–83.
15. London GM, Guerin AP, Marchais SJ, et al. Arterial media calcification in end-stage renal disease: impact on all-cause and cardiovascular mortality. *Nephrol Dial Transplant*. 2003;18(9):1731–40.
16. Gelev S, Spasovski G, Trajkovski Z, et al. Factors associated with various arterial calcifications in haemodialysis patients. *Prilozi*. 2008;29(2):185–99.
17. National Kidney Foundation. K/DOQI, NKF: clinical practice guidelines for bone metabolism and disease in chronic kidney disease. *Am J Kidney Dis*. 2003;42:S1–201.
18. Bleyer AJ, Burke SK, Dillon M, et al. A comparison of the calcium-free phosphate binder sevelamer hydrochloride with calcium acetate in the treatment of hyperphosphatemia in hemodialysis patients. *Am J Kidney Dis*. 1999;33(4):694–701.
19. Kurz P, Monier-Faugere MC, Bognar B, et al. Evidence for abnormal calcium homeostasis in patients with adynamic bone disease. *Kidney Int*. 1994;46:855–61.
20. Chertow GM, Burke SK, Raggi P. Sevelamer attenuates the progression of coronary and aortic calcification in hemodialysis patients. *Kidney Int*. 2002;62(1):245–52.
21. Zoccali C, Abramowicz D, Cannata -Andia JB, et al. European best practice quo vadis? From European best practice guidelines (EBPG) to European renal best practice (ERBP). *Nephrol Dial Transplant*. 2008;23:2162–6.
22. KDIGO clinical practice guideline for the diagnosis, evaluation, prevention, and treatment of Chronic Kidney Disease-Mineral and Bone Disorder (CKD-MBD). Kidney Disease: Improving Global Outcomes (KDIGO) CKD-MBD Work Group. *Kidney Int Suppl*. 2009;(113):S1–130.
23. Farkouh ME, Fuster V. Meta-analysis of small trials: proceed with caution. *Nat Clin Pract Nephrol*. 2008;4:115.
24. Tonelli M, Wiebe N, Culleton B, et al. Alberta Kidney Disease Network. Systematic review of the clinical efficacy and safety of sevelamer in dialysis patients. *Nephrol Dial Transplant*. 2007;22:2856–66.
25. Manns B, Klarenbach S, Lee H, Culleton B, Shrive F, Tonelli M. Economic evaluation of sevelamer in patients with end-stage renal disease. *Nephrol Dial Transplant*. 2007;22(10):2867–78.
26. Block GA, Raggi P, Bellasi A, Kooienga L, Spiegel DM. Mortality effect of coronary calcification and phosphate binder choice in incident hemodialysis patients. *Kidney Int*. 2007;71(5):438–41.
27. Suki WN, Zabaneh R, Cangiano JL, et al. Effects of sevelamer and calcium-based phosphate binders on mortality in hemodialysis patients. *Kidney Int*. 2007;72(9):1130–7.
28. St Peter WL, Liu J, Weinhandl E, Fan Q. A comparison of sevelamer and calcium-based phosphate binders on mortality, hospitalization, and morbidity in hemodialysis: a secondary analysis of the Dialysis Clinical Outcomes Revisited (DCOR) randomized trial using claims data. *Am J Kidney Dis*. 2008;51(3):445–54.
29. Ferreira A, Frazão JM, Monier-Faugere MC, et al. Effects of sevelamer hydrochloride and calcium carbonate on renal osteodystrophy in hemodialysis patients. *J Am Soc Nephrol*. 2008;19(2):405–12.
30. Assimon MM, Mousa S, Shaker O, Pai AB. The effect of sevelamer hydrochloride and Calcium—based phosphate binders on mortality in hemodialysis patients: a need for more research. *Consult Pharm*. 2010;25(1):41–54.
31. Jamal SA, Fitchett D, Lok CE, et al. The effects of calcium-based versus non-calcium-based phosphate binders on mortality among patients with chronic kidney disease: a meta-analysis. *Nephrol Dial Transplant*. 2009;24(10):3168–74.
32. Shantouf R, Ahmadi N, Flores F, et al. Impact of phosphate binder type on coronary artery calcification in hemodialysis patients. *Clin Nephrol*. 2010;74(1):12–8.
33. Brandenburg VM, Schlieper G, Heussen N, et al. Serological cardiovascular and mortality risk predictors in dialysis patients receiving sevelamer: a prospective study. *Nephrol Dial Transplant*. 2010;25(8):2672–9.
34. Zhang Q, Li M, Lu Y, et al. Meta-analysis comparing sevelamer and calcium-based phosphate binders on cardiovascular calcification in hemodialysis patients. *Nephron Clin Pract*. 2010;115(4):c259–67.
35. Fishbane S, Delmez J, Suki WN, et al. A randomized, parallel, open-label study to compare once-daily sevelamer carbonate powder dosing with thrice-daily sevelamer hydrochloride tablet dosing in CKD patients on hemodialysis. *Am J Kidney Dis*. 2010;55(2):307–15.
36. De Francisco AL, Leidig M, Covic A, et al. Evaluation of calcium acetate/magnesium carbonate as a phosphate binder compared with sevelamer hydrochloride in haemodialysis patients: a controlled randomized study (CALMAG study) assessing efficacy and tolerability. *Nephrol Dial Transplant*. 2010;25:3707–17.
37. Spasovski G, Vanholder R. Is combined calcium/magnesium phosphate binder really non-inferior to sevelamer hydrochloride? *Nephrol Dial Transplant*. In press 2011.
38. Negri AL. Phosphate binders, cardiovascular calcifications and mortality: do we need another survival study with sevelamer? *J Nephrol*. 2010;23(6):653–7.
39. Manns B, Klarenbach S, Lee H, et al. Economic evaluation of sevelamer in patients with end-stage renal disease. *Nephrol Dial Transplant*. 2007; 22:2867–78.
40. White CA, Jaffey J, Magner P. Cost of applying the K/DOQI guidelines for bone metabolism and disease to a cohort of chronic hemodialysis patients. *Kidney Int*. 2007;71:312–7.

Publish with Libertas Academica and every scientist working in your field can read your article

“I would like to say that this is the most author-friendly editing process I have experienced in over 150 publications. Thank you most sincerely.”

“The communication between your staff and me has been terrific. Whenever progress is made with the manuscript, I receive notice. Quite honestly, I’ve never had such complete communication with a journal.”

“LA is different, and hopefully represents a kind of scientific publication machinery that removes the hurdles from free flow of scientific thought.”

Your paper will be:

- Available to your entire community free of charge
- Fairly and quickly peer reviewed
- Yours! You retain copyright

<http://www.la-press.com>