

OPEN ACCESS

Full open access to this and thousands of other papers at <http://www.la-press.com>.

Cranberry (*Vaccinium macrocarpon*) changes the surface hydrophobicity and biofilm formation of *E. coli*

Jose Uberos¹, Ishan Iswaldi², Rocio Rodríguez Belmonte¹, Antonio Segura-Carretero², Verónica Fernández-Puentes¹, Antonio Molina-Carballo¹ and Antonio Muñoz-Hoyos¹

¹Department of Paediatrics. Faculty of Medicine, University of Granada. ²Department of Analytical Chemistry, Faculty of Sciences, University of Granada, Spain. Corresponding author email: uberosfernandez@terra.es

Abstract: Cranberry has been shown useful in the prevention of urinary infection by *E. coli*. In this study, we examined the changes in the hydrophobicity of P fimbriated *E. coli* and biofilm formation after incubation with commercial cranberry syrup extract at various concentrations. After incubating a bacterial suspension with cranberry at dilutions of 1:100 and 1:1000, tests of haemagglutination inhibition, surface hydrophobicity and biofilm formation were carried out. The surface hydrophobicity of *E. coli* decreased significantly after incubation with cranberry and this effect was not modified by the culture medium. Biofilm formation was inhibited after incubation with cranberry syrup and this effect was dependent on the culture medium. Thus, in some circumstances cranberry can modify nonspecific adhesive properties of *E. coli*. In previous studies, cranberry has only been implicated in the inhibition of P-fimbriated *E. coli*, but our observations show that it acts by modifying adhesive properties under P-related fimbriae.

Keywords: cranberry syrup, *E. coli*, biofilm, hydrophobicity, P-fimbriae

Microbiology Insights 2011:4 21–27

doi: [10.4137/MBI.S8526](https://doi.org/10.4137/MBI.S8526)

This article is available from <http://www.la-press.com>.

© the author(s), publisher and licensee Libertas Academica Ltd.

This is an open access article. Unrestricted non-commercial use is permitted provided the original work is properly cited.



Introduction

Cranberries contain two chemical constituents that have been associated with the prevention of bacterial adherence: fructose (a monosaccharide) and proanthocyanidins (PACs). Fructose inhibits the adherence of mannose sensitive fimbriae to urinary epithelial cells.¹ Various studies have shown that cranberry PACs inhibit the adherence of P fimbriated *E. coli* to the urinary epithelium, while having only slight effect on Type 1 fimbriae.^{1,2}

The adherence of *E. coli* to the epithelia is considered the first step in the development of infection, and so the relation between its pathogenic effect and the tissue's attachment capacity is of great importance for subsequent disease development.³ Oligosaccharide chains containing mannose are frequently found in cell surface glycoproteins, and thus many bacteria have developed fimbriae to adhere to these molecules (Type 1 fimbriae), which can be blocked when mannose is added. Together with the expression of Type 1 fimbriae, some *E. coli* strains express a specific lectin (α -Gal(1 \rightarrow 4) β -Gal) associated with P fimbriae, which mediate the specific adherence of *E. coli* to the urinary epithelial cells.¹ Both Type 1 (mannose-sensitive) and Type P (mannose-resistant) fimbriae of *E. coli* are regulated by genes that are localised in the bacterial chromosome, and their expression can be modified depending on the characteristics of the culture medium or of the host's internal medium. Some authors⁴ have suggested that *Vaccinium macrocarpon* syrup may induce conformational changes in the surface macromolecules of P fimbriae, reducing both their length and their density. Thus, the expression change of the lectins on the bacterial surface is a determinant factor, on the one hand, in the specific adherence to epithelial cell receptors and, on the other, in modifying certain physical characteristics of the bacteria (hydrophobicity) that favour or hamper its approach to the epithelial surface, and hence its adherence.⁵

Mannose-resistant fimbriae share many characteristics that have previously been related to the colonization factor antigen (CFA), and more recently with P fimbriae. Several studies^{1,6,7} have shown that *Vaccinium macrocarpon* syrup inhibits the adherence of uropathogenic *E. coli* to epithelial cells, an effect that has been related with the functionality of P fimbriae (the virulence factor) and, to a lesser

degree, with Type 1 fimbriae.¹ Bacterial attachment to a substratum is the first step in biofilm formation. The contribution of type 1 fimbriae and P to biofilm formation has been studied by several authors and seems to depend on the characteristics of the culture medium, the strain and the surface characteristics.⁸ Less information is available on the effects of cranberry on the hydrophobic characteristics of *E. coli* and their ability to form the biofilm.⁹

To date, and to the best of our knowledge, no studies have been made assessing the capacity of cranberry to modify the adherent properties of bacteria that are not related to P fimbriae. The fundamental aim of the present study was to assess whether cranberry may affect the "non-specific" adherence properties of *E. coli*. In addition, we wished to determine whether the inhibition of haemagglutination by cranberry is associated with changes in nonspecific adherence properties vs surface hydrophobicity, adherence to nitrocellulose or biofilm formation by *E. coli*.

Material and Methods

Bacterial strains, media and growth conditions

Nine strains of uropathogenic *E. coli* (695, 787, 471, 472, 593, 595, 760, 629 and 607) were used, obtained from patients with acute pyelonephritis, together with 4 strains of *E. coli* obtained from the Spanish Type Culture Collection (CECT): CECT 424 (F- thr- leu- lacY mtl- thi- ara gal ton 2 malA xyl. Resistant to phages T1, T2 and T6.); CECT 4076 (Serovar. O157:H7, originally isolated from haemorrhagic colitis); CECT 417 (SupE44(am). mutant tRNA); and CECT 743 (Serovar. O142 K86B:H6. Isolated from children with diarrhoea).

In order to enhance the activity of the Type 1 fimbriae,¹⁰ the strains were grown in TSB culture medium at 37 °C for 48 h, then centrifuged at 2000 g for 10 min. The supernatant was then discarded, and the strains suspended in PBS (pH 7.4). This washing process was performed twice. Finally, the bacterial suspension was adjusted to 10⁹ bacteria/mL (OD of 1.0 to 542 nm). To enhance the activity of the P fimbriae, the strains were incubated for 16 h on CFA agar,¹¹ extracted from the surface of the agar after washing with 5 ml PBS, and the suspension was then centrifuged at 3000 rpm for 10 min. The supernatant was then discarded, the washing process was



repeated, and finally the bacteria were suspended in PBS with 1% mannose to obtain a concentration of 10^9 bacteria/mL (OD of 1 to 542 nm). All measurements were performed in duplicate and the mean values of the two measurements were used.

Haemagglutination

The blood used was obtained from donors with blood group A Rh+, and collected in tubes with 100 μ l of trisodium citrate as anticoagulant, at 3.8%, to which 1 ml of blood was added. For the haemagglutination techniques of the mannose-sensitive strains, the blood was diluted to 1:4 with PBS. For haemagglutination of the mannose-resistant strains, the blood was diluted in PBS with methyl- α -D-mannopyranoside 0.1 mM. Over a U-bottom microwell plate, 50 μ l of bacterial suspension and 50 μ l of diluted fresh blood (1:4) were added. The haemagglutination results became visible after 45–60 min at room temperature.

Cranberry syrup: procedure

The cranberry syrup was provided by Pharmatoka Lab., (France) using a dispersion of 2.8% cranberry extract Gykacran® in glucose syrup (France) and characterised by the Department of Analytical Chemistry at the University of Granada.¹² Serial dilutions of cranberry syrup were made in PBS (pH 7.4) to 1:10 and 1:100, and the pH was adjusted to 7.0 with NaOH. 100 μ l of each dilution of the cranberry extract was incubated for 30 min at room temperature in a rotary shaker with 1 ml of the bacterial suspension (OD 1 to 542 nm), producing a final dilution of the cranberry syrup extract of approximately 1:100 and 1:1000. The final dilution of 1:1000 was chosen because this was the lowest dilution that inhibited haemagglutination. Each suspension was then centrifuged at 3000 rpm for 10 min and resuspended in an equal volume of ammonium sulphate (salt aggregation) or PBS (adherence to nitrocellulose or biofilm formation).

Ammonium sulphate aggregation test

The technique used was as described by Lindahl et al.¹³ Briefly, solutions of ammonium sulphate were prepared, with osmolarities ranging from 0.2M to 4M, using sodium phosphate as a dilutant. Taking 20 μ l of bacterial suspension, an equal volume of ammonium sulphate solution was added. The presence of aggregation was observed after 30 seconds' gentle manual

rotation, at room temperature, over a glass slide, and the lowest concentration of ammonium sulphate that produced visible aggregation was noted down. Aggregation with the 4M solution was interpreted as a hydrophobicity of 0%, while aggregation with the 0.2M solution was interpreted as 95% hydrophobicity.

Adherence to nitrocellulose filters

The procedure described by Lachica and Zink¹⁴ was performed. Briefly, a bacterial suspension in a PUM buffer with ammonium sulphate 0.2M (1:2, v:v) was adjusted to 10^5 bacteria (OD 0.6 to 540 nm). The initial absorbance (Abs_1) was determined, and 3 ml of this bacterial suspension was passed through an 8 μ m-pore nitrocellulose filter (Millipore). The absorbance of the filtrate (Abs_2) was then determined and the results expressed as a percentage of the adherence to the nitrocellulose filter.

Biofilm formation¹⁵

The strains were incubated at 37 °C for 24 h in glass tubes with 2.5 ml of TSB or CFA broth. 0.5 ml of the latter and 50 μ l of cranberry syrup at dilutions 1:10 and 1:100 were placed in Eppendorf tubes, resulting in final cranberry syrup dilutions of 1:100 and 1:1000. An Eppendorf tube containing an equal amount of TSB or CFA broth without inoculum was used as a negative control, and 0.5 ml of the bacterial suspension in Eppendorf tubes as a positive control. After incubation for 24 hours, the contents of each tube were aspirated carefully with a Pasteur pipette, washed three times with 250 μ l of PBS, and gently shaken to remove non adherent bacteria. The tubes were then air dried and added to 200 μ l of 99% methanol as a fixative, and left for 15 minutes. The excess methanol was then discarded and the tubes air dried. 200 μ l of 2% Hucker's crystal violet dye was added and the tubes submerged in distilled water for 5 minutes, with the remaining tubes being discarded. This procedure was repeated three times. Subsequently, bacteria were solubilized in each tube, in 250 μ l of 33% acetic acid. The optical density of the suspension of each tube was measured at 540 nm and the zero adjusted to 33% acetic acid. Non-inoculated TSB broth was used as a negative control. The results obtained are expressed as follows: $OD \leq OD$ control: Non adherent bacteria. OD control $\leq OD \leq 2 \times OD$ control: slightly adherent bacteria. $2 \times OD$ control $\leq OD \leq 4 \times OD$ control:



moderately adherent bacteria. $OD \leq 4 \times OD$ control: Highly adherent bacteria.

Statistical analysis

SPSS 15.0 statistical software was used to perform the Friedman and Wilcoxon test for the comparison of dependent variables, and the Mann-Whitney-Wilcoxon test for the comparison of independent variables. Kendall's W was used for the comparison of paired variables. In addition, Kendall's coefficient of inter-variable concordance was calculated.

Results

Table 1 shows that the addition of methyl- α -D-mannopyranoside 0.1 mM to blood group A, Rh+ does not inhibit the haemagglutination of any of the 13 strains of *E. coli* utilised in this study (mannose-resistant strains). The incubation of strains of *E. coli* with cranberry syrup extract at a final concentration of 1:1000 did not inhibit haemagglutination, while incubation with the cranberry at a final concentration of 1:100 inhibited it, in every case, for strains grown either in TSB or CFA culture medium.

No differences were observed in the surface hydrophobicity of *E. coli* following its growth in TSB culture medium ($Z = 0.35$; pNS). However, incubation of the bacterial suspension with cranberry syrup at a final concentration of 1:1000 or 1:100 produced significant reductions in surface hydrophobicity,

depending on the concentration of cranberry, after growth in either TSB or CFA medium (Fig. 1).

The percentage of adherence to nitrocellulose was not significantly modified after growth on CFA agar or in TSB culture medium ($Z = 0.73$; pNS). In both culture media, the percentage of adherence to nitrocellulose of *E. coli*, after incubation for 20 min with cranberry syrup, remained unchanged. Kendall's test of concordance revealed the existence of concordance between the surface hydrophobicity values of *E. coli* and the percentage of adherence to nitrocellulose following growth in TSB culture medium ($T = 0.50$; $P = 0.02$), and an absence of such concordance following growth on CFA agar ($T = 0.09$; $P = NS$).

After culture in CFA broth, biofilm formation was unchanged by incubation with cranberry syrup at a final concentration of 1:1000 or 1:100 ($W = 0.138$, $P = NS$). However, in TSB medium, biofilm formation was inhibited after incubation with cranberry syrup ($W = 0.93$, $P < 0.001$), and the formation of biofilm in the absence of cranberry syrup was significantly higher after culture in TSB ($Z = 2.31$, $P = 0.02$), as shown in Figure 2.

Discussion

Our study shows that cranberry syrup can modify the surface hydrophobicity and the biofilm formation of *E. coli* on polystyrene (microtitre plates), the latter effect being dependent on the characteristics of the

Table 1. Results of the haemagglutination test (A Rh+) for mannose resistant *E. coli*. Cranberry 1:100 inhibits haemagglutination in both culture media.

<i>E. coli</i> strain	Haemagglutination TSB culture medium			Haemagglutination CFA culture medium		
	Control	Cranberry 1:1000	Cranberry 1:100	Control	Cranberry 1:1000	Cranberry 1:100
471	+	+	-	+	+	-
787	+	+	-	+	+	-
593	+	+	-	+	+	-
472	+	+	-	+	+	-
595	+	+	-	+	+	-
760	+	+	-	+	+	-
695	+	+	-	+	+	-
743	+	+	-	+	+	-
424	+	+	-	+	+	-
607	+	+	-	+	+	-
417	+	+	-	+	+	-
4076	+	+	-	+	+	-
629	+	+	-	+	+	-




Figure 1. Mean values and standard error of the mean of surface hydrophobicity for *E. coli* after incubation in CFA and TSB broth. There is a statistically significant decrease in surface hydrophobicity after incubation with cranberry syrup at final dilutions of 1:1000 and 1:100.

Note: *** $P < 0.001$.




Figure 2. Mean values and standard error of biofilm formation rates for *E. coli* after incubation in CFA and TSB broth. Growth in TSB produces a statistically significant decrease in adherence after incubation with cranberry syrup at final dilutions of 1:1000 and 1:100.

Note: * $P < 0.05$; *** $P < 0.001$.

culture medium. The greater expression of P fimbriae, as a response to the culture medium, was not found to be related with the greater formation of biofilm. The first step in the colonisation of the epithelium by *E. coli* is determined by its capacity to adhere to the host cells; this capacity is determined, initially, by the micro-organism's electric surface charge and its surface hydrophobicity,¹⁶ and subsequently by other factors, such as the formation of diverse types of fimbriae, and of specific adhesins.¹⁷⁻¹⁹ Miyazaki et al²⁰ showed that fimbriae types 1, P and S are not essential factors in the adhesion and later colonisation of the urogenital epithelium by *E. coli*. These adhesins may, in general, be considered features of the virulence of extra-intestinal *E. coli*, but they are not essential for *E. coli* to become uropathogenic, which accounts for current research interest into the extent to which cranberry extract may affect the non-specific adherence properties of *E. coli*. Cross-talk among *pilus operons* within a bacterial cell, probably triggered by environmental cues, can result in a switch in expression from one pilus type to another, a process known as phase variation. The variable

expression of the fimbriae in a single strain of *E. coli* increases the probability of adherence and invasion of the uroepithelium.²¹

Cranberries contain different amounts of fructose, PACs, flavonoids and anthocyanins. PACs isolated from cranberry fruit consist of predominantly epicatechin units with at least one A-type linkage. The A-linkage in cranberry PACs may represent an important structural feature for bacterial anti-adherence activity. The type A cranberry PACs prevent adherence of P-fimbriated uropathogenic *E. coli* to uroepithelial cells *in vitro*.⁴ The cultivation of *E. coli* in CFA broth favours the expression of P fimbriae.¹¹ We observed that the incubation of *E. coli* in CFA broth with cranberry syrup does not alter the formation of biofilm, from which we infer that the increased expression of P fimbriae is not related to the formation of biofilm. The latter finding differs from that observed after culture in TSB broth, implying that biofilm formation can be inhibited by cranberry syrup, depending on the characteristics of the culture medium, while the expression of P fimbriae is not an important element in the formation of biofilm. Hancock et al⁸ showed that




Figure 3. Mean values of adherence to nitrocellulose for *E. coli* before and after incubation with cranberry syrup at final dilutions of 1:1000 and 1:100.

the expression of P fimbriae significantly decreases biofilm formation under different growth conditions.

The incubation of *E. coli* with cranberry syrup extract produces a reduction in the surface hydrophobicity, dependent on the concentration of the extract, and which is independent of the different culture media (Fig. 1). This effect may be different from the inhibition of P fimbriae of cranberry PACs, since according to our findings and those of Hancock et al,⁸ P fimbriae have little influence on the formation of biofilm. Further studies are currently being conducted, involving a partitioning of the components of the cranberry syrup and a study of the effects thus produced. With respect to surface hydrophobicity, the incubation of *E. coli* with a 1:1000 dilution of cranberry extract, which does not affect the haemagglutination mediated by Type P fimbriae, does produce significant reductions in the surface hydrophobicity, thus showing that extremely low levels of this extract are capable of modifying the non-specific adherence properties of *E. coli*. According to our own results, the adherence of *E. coli* to nitrocellulose is not related to the predominant expression of Type P or Type 1

fimbriae, and incubation with cranberry syrup extract does not affect adherence rates.

We conclude that, depending on the culture medium, cranberry can modify the non-specific adherence properties of *E. coli* and alter its ability to form biofilm.

Abbreviations

TSB, Tryptic Soy Broth; CFA, Colonization Factor Antigen; PBS, Phosphate buffered saline.

Acknowledgements

This study was funded by the Carlos III Institute of Health (Spain).

Disclosures

Author(s) have provided signed confirmations to the publisher of their compliance with all applicable legal and ethical obligations in respect to declaration of conflicts of interest, funding, authorship and contributorship, and compliance with ethical requirements in respect to treatment of human and animal test subjects. If this article contains identifiable human subject(s) author(s) were required to supply signed patient consent prior to publication. Author(s) have confirmed that the published article is unique and not under consideration nor published by any other publication and that they have consent to reproduce any copyrighted material. The peer reviewers declared no conflicts of interest.

References

- Zafiri D, Ofek I, Adar R, Pocino M, Sharon N. Inhibitory activity of cranberry juice on adherence of type 1 and type P fimbriated Escherichia coli to eucaryotic cells. *Antimicrob Agents Chemother.* 1989;33:92–8.
- Foo LY, Lu Y, Howell AB, Vorsa N. The structure of cranberry proanthocyanidins which inhibit adherence of uropathogenic P-fimbriated Escherichia coli in vitro. *Phytochemistry.* 2000b;54:173–81.
- Sellwood R, Gibbons RA, Jones GW, Rutter JM. Adhesion of enteropathogenic Escherichia coli to pig intestinal brush borders: the existence of two pig phenotypes. *J Med Microbiol.* 1975;8:405–11.
- Howell AB. Bioactive compounds in cranberries and their role in prevention of urinary tract infections. *Mol Nutr Food Res.* 2007;51:732–7.
- Magnusson KE. Hydrophobic interaction--a mechanism of bacterial binding. *Scand J Infect Dis Suppl.* 1982;33:32–6.
- Foo LY, Lu Y, Howell AB, Vorsa N. A-Type proanthocyanidin trimers from cranberry that inhibit adherence of uropathogenic P-fimbriated Escherichia coli. *J Nat Prod.* 2000a;63:1225–8.
- Foo LY, Lu Y, Howell AB, Vorsa N. The structure of cranberry proanthocyanidins which inhibit adherence of uropathogenic P-fimbriated Escherichia coli in vitro. *Phytochemistry.* 2000c;54:173–81.
- Hancock V, Witsø IL, Klemm P. Biofilm formation as a function of adhesin, growth medium, substratum and strain type. *Int J Med Microbiol.* 2011;301:570–6.



9. Liu Y, Black MA, Caron L, Camesano TA. Role of cranberry juice on molecular-scale surface characteristics and adhesion behavior of *Escherichia coli*. *Biotechnol Bioeng*. 2006;93:297–305.
10. Blanco M, Blanco J, Blanco JE, et al. Virulence factors and O serogroups of *Escherichia coli* as a cause of community-acquired urinary infections. *Enferm Infecc Microbiol Clin*. 1995;13:236–41.
11. Evans DG, Evans DJ Jr, Tjoa W. Hemagglutination of human group A erythrocytes by enterotoxigenic *Escherichia coli* isolated from adults with diarrhea: correlation with colonization factor. *Infect Immun*. 1977;18:330–7.
12. Iswaldi I, Gómez-Caravaca AM, Uberos J, Lardón M, Segura-Carretero A, Fernández-Gutierrez A. Characterization by high-performance liquid chromatography with diode-array detection coupled to time-of-flight mass spectrometry of the phenolic fraction in a cranberry syrup used to prevent urinary tract diseases, together with a study of its antibacterial activity. *J Pharm Biomed Anal*. In press. 2011.
13. Lindahl M, Faris A, Wadstrom T, Hjerten S. A new test based on ‘salting out’ to measure relative surface hydrophobicity of bacterial cells. *Biochim Biophys Acta*. 1981;677:471–6.
14. Lachica RV, Zink DL. Plasmid-associated cell surface charge and hydrophobicity of *Yersinia enterocolitica*. *Infect Immun*. 1984;44:540–3.
15. Stepanovic S, Vukovic D, Dakic I, Savic B, Svabic-Vlahovic M. A modified microtiter-plate test for quantification of staphylococcal biofilm formation. *J Microbiol Methods*. 2000;40:175–9.
16. Ferreirós CM, Criado MT. Expression of surface hydrophobicity encoded by R-plasmids in *Escherichia coli* laboratory strains. *Arch Microbiol*. 1984;138:191–4.
17. Ferreirós CM, Criado MT. Adhesive properties associated with the K99 antigen of *Escherichia coli*. *IRCS Medical Science*. 1980;10:917–8.
18. Gibbons RJ, Spinell DM, Skobe Z. Selective adherence as a determinant of the host tropisms of certain indigenous and pathogenic bacteria. *Infect Immun*. 1976;13:238–46.
19. Gibbons RJ, Hay DL. Adsorbed salivary acidic proline-rich proteins contribute to the adhesion of *Streptococcus mutans* JBP to apatitic surfaces. *J Dent Res*. 1989;68:1303–7.
20. Miyazaki J, Ba-Thein W, Kumao T, Obata YM, Akaza H, Hayashi H. Type 1, P and S fimbriae, and afimbrial adhesin I are not essential for uropathogenic *Escherichia coli* to adhere to and invade bladder epithelial cells. *FEMS Immunol Med Microbiol*. 2002;33:23–6.
21. Wiles TJ, Kulesus RR, Mulvey MA. Origins and virulence mechanisms of uropathogenic *Escherichia coli*. *Exp Mol Pathol*. 2008;85:11–9.

Publish with Libertas Academica and every scientist working in your field can read your article

“I would like to say that this is the most author-friendly editing process I have experienced in over 150 publications. Thank you most sincerely.”

“The communication between your staff and me has been terrific. Whenever progress is made with the manuscript, I receive notice. Quite honestly, I’ve never had such complete communication with a journal.”

“LA is different, and hopefully represents a kind of scientific publication machinery that removes the hurdles from free flow of scientific thought.”

Your paper will be:

- Available to your entire community free of charge
- Fairly and quickly peer reviewed
- Yours! You retain copyright

<http://www.la-press.com>