

ORIGINAL RESEARCH

OPEN ACCESS
Full open access to this and
thousands of other papers at
<http://www.la-press.com>.

Temporal Variations of Atmospheric CO₂ in Dehradun, India during 2009

Neerja Sharma¹, Rabindra K Nayak¹, Vinay K Dadhwal¹, Yogesh Kant² and Meer M Ali¹

¹National Remote Sensing Centre, Hyderabad. ²Indian Institute of Remote Sensing, Dehradun.
Corresponding author email: neerja_s@rediffmail.com

Abstract: The present study reports the temporal variations of CO₂ mixing ratio measured using Vaisala GMP-343 sensor (at 15 m height) in Dehradun (30.1 °N, 77.4 °E) during 2009. Being a valley station, the mixing ratios are controlled by biospheric processes but not by large scale transport phenomenon or local pollution. A distinct diurnal cycle varies from 317.9 ppm in the afternoon to 377.2 ppm in the morning (before sunrise). The minimum early morning (0700–1000 IST) drop and minimum afternoon (1300–1700 IST) trough observed during monsoon months are related to the enhanced vegetation activity due to rain at the site. The maximum night time (2200 IST to next day 0700 IST) build up of CO₂ observed during monsoon season is associated with the increase in heterotrophic respiration due to high moisture content in the soil. This is also confirmed by the positive coherence between night time CO₂ mixing ratio with soil respiration simulated from Carnegie-Ames-Stanford Approach (CASA) model. The strong negative coherence with net ecosystem productivity (simulated from the same model) shows that observations captured the regional changes in emission and uptake of CO₂ in atmosphere.

Keywords: carbon dioxide, photosynthesis, soil respiration, ecosystem model

Air, Soil and Water Research 2013:6 37–45

doi: [10.4137/ASWR.S10590](https://doi.org/10.4137/ASWR.S10590)

This article is available from <http://www.la-press.com>.

© the author(s), publisher and licensee Libertas Academica Ltd.

This is an open access article. Unrestricted non-commercial use is permitted provided the original work is properly cited.



Introduction

Carbon dioxide (CO₂) is the most important greenhouse gas present in the atmosphere. It accumulates in the atmosphere and plays an important role in controlling the planet's surface temperature. It can be measured using sensors and satellite observations. For example, the Global Atmosphere Watch (GAW) Programme of the World Meteorological Organization (WMO) focuses on the long term greenhouse gas monitoring network to understand the climate related changes.¹ However, the complete understanding of CO₂ source and sink requires multiple approaches (with tower measurements, ecosystem analysis, and modeling etc.) due to its complexity of the interactions at different time and space scales. Atmospheric CO₂ measurement is the best approach to study the CO₂ variability on a diurnal and spatial scale.

CO₂ mixing ratio has been measured at (a) long term monitoring station, away from anthropogenic activity,² (b) tall towers where altitudinal variations permit source and sink analysis, (c) near the surface using flux tower, and (d) over urban and complex human affected regions to infer the source and sink pattern and variability. The examples of such studies are listed below. Idso et al^{3,4} used transects of Phoenix, Arizona, at 2 m to infer the presence of an urban CO₂ "dome." A similar experiment was conducted in Rome based on low level (2 m) measurements.⁴ The seasonal CO₂ mixing ratios were also analyzed in several places including 40 m tower measurements taken in suburban Melbourne, Australia.⁶ Significant diurnal cycles in CO₂ surface flux and mixing ratios have been found in Vancouver.⁷ Mixing ratio measurement from flux tower using eddy covariance method has been carried out in Copenhagen.^{6,8} Rigby et al⁹ reported the diurnal and seasonal CO₂ mixing ratios measured at 87 m tall tower in central London.

In India, Tiwari et al¹⁰ studied atmospheric CO₂ observations as a separate tracer of terrestrial, oceanic fluxes and fossil fuel emissions at coastal station Cape Rama. Bhattacharya et al¹¹ infer the relationship of south west monsoon and CO₂ mixing ratio for the same station measured from 1993 to 2002. Schuck et al¹² identified the summer monsoon signature on high altitude CO₂ mixing ratio measured by CARBIC aircraft between Germany and India during 2008. Recently, measurement and scaling of CO₂ exchanges in wheat using flux-tower was carried out by Patel et al.¹³

Though in situ observations are important and critical, very few studies have been carried out over India. A continuous measurement provides diurnal pattern which could be decomposed into day time net exchange and night time respiration release. As part of a comprehensive approach to carbon cycle (C-cycle) observations and modeling, National Carbon Project (NCP) of Indian Space Research Organization-Geosphere Biosphere Programme has initiated 3 types of CO₂ observations: (i) eddy flux, (ii) continuous atmospheric observations, and (iii) satellite derivation. Continuous measurements of atmospheric CO₂ is carried out using Vaisala GMP-343 sensor in Dehradun, Uttarakhand, the first site established during October 2008. The numerous studies which have used this sensor reported reliable results in atmospheric,^{9,14} subterranean ventilation,¹⁵ selection of space plant growth,¹⁶ variations within a canopy,¹⁷ limnology and oceanographic studies¹⁸ and in soil measurements.¹⁹

In this paper, we analyzed temporal variations in atmospheric CO₂ mixing ratios in Dehradun site during 2009 and relate them to estimated exchange by biospheric model. However, the impact of long range transport is studied using back trajectory analysis and impact of local pollution by separating mixing ratios during working and non-working days.

Site and Instrument

Dehradun (30.1 °N, 77.4 °E) is a valley, located in the Shivalik range of the Himalayas at a mean altitude of 700 m. The city has a large area covered under vegetations and trees. The climate of the city varies from hot in summer to severely cold in winter and rain during monsoon season. The Vaisala GMP-343 sensor was installed at the top of a building approximately 15 m from the ground, in the campus of the Indian Institute of Remote Sensing (IIRS). It is mounted sufficiently above and away from the roof top to avoid heating effect of the building on CO₂ measurements. Since the building is located on institute premises, which is away from the main traffic, the impact of public vehicular pollution is expected to be negligible.

CO₂ observations at 15 minute intervals were taken from the sensor through a data logger. It is a diffusion type instrument where the air diffuses into the measurement chamber through a filter without the requirement of the air to be pumped. It also provides



an acceptable compromise between size, response time accuracy, and stability. The technical specifications of the sensor and the sensitivity of the instrument to temperature, pressure, and relative humidity are given in Table 1.²⁰

Data and Methodologies

Data

In addition to CO₂ observations, the following ancillary data was also acquired:

- a. Meteorological parameters: 3 hourly temperature (°C), pressure (hPa), and relative humidity (RH) data from India Meteorological Department (IMD), Dehradun.
- b. Parameters used for running the Carnegie-Ames-Stanford Approach (CASA) model: The estimates of Net Ecosystem Productivity (NEP) and respiration were computed using the CASA model for which the inputs and their sources are:
 - i. Normalized Vegetation Index (NDVI) data were based on Moderate Resolution Imaging Spectroradiometer (MODIS) available at 1 km spatial grid over the globe;
 - ii. Data of climatic parameters are part of past 100 year data base of high resolution (0.5 × 0.5 °) temperature, precipitation and cloudiness prepared by Climate Research Unit (CRU) at the University of East Anglia (UEA) (<http://www.cru.uea.ac.uk/cru/data/hrg>). The CRU cloud fraction data are used to obtain surface solar radiation over the country at monthly time scales by using the methodology described by Mani;²¹

- iii. Extra-terrestrial radiation is obtained from the Surface Radiation Budget (SRB) databases (<http://gewex-srb.larc.nasa.gov/>);
- iv. The land cover map of the country is based on the land cover map of Southeast Asia derived at a 1 km spatial resolution by using multi-date SPOT-VEGETATION data for the global land cover mapping project;²² and
- v. The soil attribute map of the country obtained from Food and Agriculture Organization (FAO) of UNESCO's world soil map.²³

Methodology

The sensor was installed at Dehradun during October 2008 after factory calibration by Vaisala at 25 °C and 1013 hPa. Since Dehradun is located at 700 m, CO₂ measurements made at this elevation have to be corrected for the ambient temperature and pressure values. For any Non Dispersive Infra Red sensor these corrections are based on the ideal gas law, therefore we used equation (i) in the present study.²⁰

$$CO_{2corrected} \text{ (ppm)} = CO_{2measured} \text{ (ppm)} * \left(\frac{T_{measured}}{T_{reference}} \right)^* \left(\frac{P_{reference}}{P_{measured}} \right) \quad (1)$$

where,

$P_{measured}$ = Measured atmospheric pressure (hPa)

$T_{measured}$ = Measured atmospheric temperature (K)

$P_{reference}$ = Reference barometric pressure (1013 hPa)

$T_{reference}$ = Reference temperature (298 K)

The hourly averages are obtained from corrected values at 15 minute intervals.

A preliminary analysis indicated the diurnal CO₂ mixing ratios clearly demarcated (a) early morning decline, (b) afternoon low followed continuing into the evening, and (c) slow increase over night time. Thus, for seasonal behavior analysis CO₂ mixing ratio variations from 0700 IST to 1000 IST (period-I), 1200 IST to 1700 IST (period-II) and from 2200 IST to next day 0700 IST (period-III) were analyzed separately.

To understand the influence of carbon exchange on monthly variations, CASA model simulated soil respiration and NEP have been analyzed. The influence of vehicular pollution on measurements was examined

Table 1. Vaisala GMP343 key performance statistics and effect of temperature, pressure, relative humidity on sensor performance.

Accuracy	At the CO ₂ calibration points: ±1.5% of reading
Stability	<±2% of reading/year
Response time	30 sec.
Noise at 350 ppm	±3 ppm
Temperature accuracy (% of reading) (-10 °C to +40 °C)	±0.5
Pressure (900 to 1050 hPa)	±0.5
Relative humidity	±0.006

by evaluating the difference of CO₂ values on working and non working days. The effect of large scale transport is evaluated using back trajectory analysis.^{24,25}

The CASA Model

The CASA model²⁶ is a biogeochemical model, which simulates the cycling of carbon between plants, soil, and atmosphere as the exchange of carbon between biogeochemical pools. Calculation of monthly terrestrial Net Primary Productivity (NPP) is based on the concept of light-use efficiency modified by temperature and moisture stress. Soil carbon cycling and heterotrophic respiration (Rh) flux components of the model are based on a compartmental pool structure, with first-order equations to simulate loss of CO₂ from decomposing plant residue and surface soil organic matter (SOM). NEP, which is the difference between NPP and Rh, determines the estimation of net carbon exchange between the terrestrial ecosystems. The model has been implemented by several workers to estimate regional or continental patterns of NPP and the CO₂ sink over North America, Eurasia, South America, Africa, and Australia.^{27–29}

Recently, Nayak et al^{30,31} implemented this model for the Indian terrestrial ecosystem. For this purpose, monthly climatic drivers and NDVI are sampled at 2 min spatial resolution in order to match the model grid size. The soil map provides the data of various compositions of the soil in the upper 30 cm depth (sand, silt, and clay fraction) at a 5 min resolution. Nayak et al³¹ interpolated this data over the subcontinent at the model resolution and then a soil attribute map, as per the seven classes defined in the CASA model. The model is forced by time varying climate data bases, NDVI, and stationary maps of soil and vegetation to simulate monthly carbon parameters.

Results

Variations in the meteorological parameters

The monthly variations of meteorological parameters: (a) relative humidity (%), (b) temperature (°C), and (c) pressure (hPa) in Dehradun during 2009 are shown in Figure 1. Summer at this station is generally from March to June, monsoon months from July to October and winter from November to February. The winter is cold (temperature is less than 20 °C) and humid (RH is above 60%) with high atmospheric pressure (above 930 hPa).




Figure 1. Monthly variability of atmospheric parameters (A) relative humidity (%), (B) temperature (°C), and (C) pressure (hPa) in Dehradun during 2009.

Temperature increases from 16 °C in February to 31 °C in June with low humid conditions (below 50%). From July to October temperature remains above 24 °C with high humid conditions (above 70%) and with low atmospheric pressure. The monthly variations in atmospheric pressure shows that the atmospheric boundary layer is dense during winter and rare during summer months, which can strongly contribute to CO₂ mixing ratio variations as explained later.

CO₂ analysis

Monthly average diurnal cycles

Monthly average hourly CO₂ mixing ratios exhibit a clear diurnal cycle (Fig. 2). The maximum (377.1 ppm) occurs in morning and minimum (317.9 ppm) in afternoon. The rapid decrease from morning maximum is mainly due to atmospheric mixing, whereas the gradual afternoon drop is because of net ecosystem uptake. The night time build up is mainly governed by increases in boundary layer stability and cessation of photosynthesis activity. Similarly, near surface features of CO₂ variability is also reported earlier.^{32,33} Though the response of diurnal cycles based on these processes are similar, they have large monthly variations.

Analysis during three periods of the day

Period I: The monthly average difference of CO₂ mixing ratios from 0700 to 1000 IST is shown in




Figure 2. Diurnal cycles of hourly CO₂ mixing ratio in Dehradun from January to December during 2009.

Figure 3. The difference starts increasing from January (9.7 ppm) and reaches the maximum (35.0 ppm) during July–October and decreases afterwards. Thus, the magnitude of the morning drop is low during winter and pre-monsoon when compared to monsoon and post-monsoon drops. Respired CO₂ accumulated more over the vegetated area near dawn during growing season. With an increase in photosynthesis activity, the accumulated mixing ratio decreases at a faster rate therefore the drop is at its highest during monsoon and post-monsoon seasons. Spittlehouse




Figure 3. Difference of monthly average CO₂ mixing ratio between 0700 and 1000 IST.

and Ripley³⁴ also reported the maximum morning drop over a vegetated area during growing seasons. On the contrary, with the winter and pre-monsoon seasons being less productive, the drop in CO₂ is less pronounced.

Period II: Monthly average CO₂ mixing ratio from 1300 to 1700 IST is shown in Figure 4. It varies from 317.9 ppm in September to 336.0 ppm in December. The pattern shows bi-annual variability with two crests (during December/January and April) and two troughs (during March and September). From January (331.7 ppm) it starts decreasing and reaches to the first minimum in March (328.9 ppm), and then increases until June. Then it decreases and reaches the minimum in September from where it increases once again. Day time uptake of CO₂ is high during monsoon and post-monsoon seasons due to increases in vegetation activity. The opposite can be said about daytime uptake in the winter and pre-monsoon seasons. Uptake of CO₂ by vegetation leads to a low mixing ratio observed in London city as explained by Rigby et al.⁹

Period III: The hourly CO₂ mixing ratio from 2200 IST to next day of 0700 IST during winter, pre-monsoon, monsoon and post-monsoon seasons are shown in Figure 5. The night time build up of CO₂ during this period is the result of three processes: (i) absence of photosynthesis activity, (ii) stable atmospheric boundary layer, and (iii) continuous emission due to plants and soil.³⁵ The continuous high mixing ratios are observed during winter nights, whereas it increases from night to morning during




Figure 4. Monthly average CO₂ mixing ratio from 1300 to 1700 IST.




Figure 5. Hourly averages of CO₂ mixing ratio from 2200 to 0700 IST during winter, pre-monsoon, monsoon and post-monsoon seasons.

other seasons. The high pressure during winter nights (Fig. 1) generally forms a nocturnal boundary layer that accumulates the pollutants within that height.³⁶ Therefore high mixing ratios are observed throughout the night. During pre-monsoon, monsoon and post-monsoon seasons, the increase in mixing ratios are 4.4 ppm, 8.2 ppm and 7 ppm respectively. The maximum increase observed during monsoon season may be attributed to high moisture content in the soil due to rains.

Comparison with model results

To further explore the causes of monthly night time and total monthly variations of CO₂, we compared our observations with the independent CASA model simulated soil respiration and NEP. The monthly soil respiration derived from the CASA model follows the average monthly night time CO₂ mixing ratio throughout the year (Fig. 6). High mixing ratio and high soil respiration are observed during monsoon months. This confirms that more moisture in the soil increases the heterotrophic respiration. To evaluate the response of total local ecosystem activity, the monthly average mixing ratios are compared with the CASA model simulated NEP (Fig. 7). A negative NEP value means NPP is less than respiration; hence CO₂ is released to the atmosphere. Whereas, when NPP is exceeding soil respiration (a positive NEP), CO₂ is being taken in by the ecosystem from the atmosphere. Therefore, both show strong negative coherence throughout the year. The low NEP




Figure 6. Comparison between night time (from 2200 IST to 0700 IST) CO₂ and CASA model simulated soil-respiration during 2009.

and high mixing ratios are observed during winter season. Enhanced ecosystem productivity decreases the CO₂ during monsoon season.

Long range transport

In India, the large point sources (LPS) associated with fossil fuel combustion and agriculture sectors are mainly located in the Mumbai-Ahmedabad corridor, Delhi, and near the coal mine mouth.³⁷ To explore the influence of air masses coming from these LPS on CO₂, we analyzed initial 7 day back trajectories for all the months (Fig. 8). Flow pattern during different months shows that Dehradun is mainly influenced by the air masses coming from the northern regions during winter, pre-monsoon and post-monsoon seasons and, as expected, from the southwest direction during




Figure 7. Comparison between monthly average CO₂ and CASA model simulated NEP during 2009.




Figure 8. First 7 days back trajectories of months from January to December during 2009.

monsoon season. Though CO₂ is a well mixed gas, from these trajectories we can infer that observation site is not influenced by the air masses coming from LPS. Therefore, the impact of long range transport on CO₂ mixing ratios at this station is ruled out.

Impact of local vehicular pollution

Since, the measurement site is away from the main traffic area, we examined the impact of local vehicle movement inside the campus on CO₂ mixing ratio. For this purpose, we computed the annual average mixing ratios between working and non-working days. Since the difference between the working (349.3 ppm) and non-working (348.9 ppm) days is negligible the effect of internal vehicular pollution is considered insignificant.

Conclusions

15 min CO₂ observations from Vaisal GMP-343 are averaged on hourly basis after applying temperature and pressure corrections at Dehradun during 2009.

Monthly average hourly mixing ratios have a clear diurnal cycle with morning maximum, afternoon minimum, and night time rise. These are mainly controlled by the photosynthesis activity, boundary layer stability, and night time respiration. Though the response of diurnal cycles based on these processes are similar, they have large monthly variations. The large decline from morning maximum observed during monsoon and post-monsoon seasons is associated with the accumulation of CO₂ on vegetated areas near dawn. Similarly, high vegetation activity during monsoon and post-monsoon seasons decreases afternoon mixing ratios more effectively. The continuously high mixing ratios during winter nights are associated with the stable boundary layer. However, the maximum night time increase observed during monsoon is due to enhanced heterotrophic respiration because of high soil moisture. CASA model simulated NEP being in opposite phase with CO₂ mixing ratio throughout the year which indicates the dominant role played by



the biospheric processes in controlling the carbon exchange. Back trajectory analysis and the amount of mixing ratio between working and non-working days further confirm that the biospheric processes exert a strong control on temporal variations of CO₂ mixing ratios at the site.

Acknowledgements

This research work was carried out as the part of National carbon Project, ISRO-Geosphere and Biosphere Programme. The authors are thankful to the referees for their critical comments and suggestions which help to improve the clarity of the manuscript.

Author Contributions

Conceived and designed the experiments: VKD. Analysed the data: NS, RKN, YK. Wrote the first draft of the manuscript: NS, MMA, VKD. Contributed to the writing of the manuscript: NS, MMA, RKN, VKD. Agree with manuscript results and conclusions: All authors. Jointly developed the structure and arguments for the paper: VKD, NS, MMA. Made critical revisions and approved final version: VKD. All authors reviewed and approved of the final manuscript.

Funding

This project is funded by the National Carbon Project (NCP) of ISRO.

Competing Interests

Author(s) disclose no potential conflicts of interest.

Disclosures and Ethics

As a requirement of publication author(s) have provided to the publisher signed confirmation of compliance with legal and ethical obligations including but not limited to the following: authorship and contribution, conflicts of interest, privacy and confidentiality and (where applicable) protection of human and animal research subjects. The authors have read and confirmed their agreement with the ICMJE authorship and conflict of interest criteria. The authors have also confirmed that this article is unique and not under consideration or published in any other publication, and that they have permission from rights holders to reproduce any copyrighted material. Any disclosures

are made in this section. The external blind peer reviewers report no conflicts of interest.

References

1. WMO Global Atmosphere Watch (GAW) Strategic Plan: 2008–2011. WMO TD No. 1384.
2. Keeling CD. The concentration and isotopic abundances of carbon dioxide in the atmosphere. *Tellus*. 1960;12(2):200–3.
3. Idso CD, Idso SB, Balling RC. The urban CO₂ dome of Phoenix, Arizona. *Phys Geo*. 1998;19:95–108.
4. Idso CD, Idso SB, Balling RC. An intensive two-week study of an urban CO₂ dome in Phoenix, Arizona, USA. *Atmos Environ*. 2001;35:995–1000.
5. Gratani L, Varone L. Daily and seasonal variation of CO₂ in the city of Rome in relationship with the traffic volume. *Atmos Environ*. 2005;39:2619–24.
6. Coutts A, Beringer M, Tapper J. Characteristics influencing the variability of urban CO₂ fluxes in Melbourne, Australia. *Atmos Environ*. 2007;41(1): 51–62.
7. Reid KH, Steyn DG. Diurnal variations of boundary-layer carbon dioxide in a coastal city—observations and comparison with model results. *Atmos Environ*. 1997;31(18):3101–14.
8. Soegaard H, Moller-Jensen L. Towards a spatial CO₂ budget of a metropolitan region based on textural image classification and flux measurements. *Remote Sens Environ*. 2003;87(2–3):283–94.
9. Rigby M, Toumi R, Fisher R, Lowry D, Nisbet EG. First continuous measurements of CO₂ mixing ratio in central London using a compact diffusion probe. *Atmos Environ*. 2008;42:8943–53.
10. Tiwari YK, Patra PK, Chevallier F, et al. Carbon dioxide observations at Cape Rama, India for the period 1993–2002 implications for constraining Indian emissions. *India Curr Sci*. 2011;101:1562–8.
11. Bhattacharya SK, Borole DV, Francey RJ, et al. Trace gases and CO₂ isotope records from Cabo de Rama. *India Curr Sci*. 2009;97:1336–44.
12. Schuck TJ, Brenninkmeijer CAM, Baker AK, Slemr F, von Velthoven PFJ, Zahn A. Greenhouse gas relationships in the Indian summer monsoon plume measured by the CARIBIC passenger aircraft. *Atmos Chem Phys*. 2010;10:3965–84.
13. Patel, NR, Dadhwal VK, Shah SK. Measurement and Scaling of Carbon Dioxide (CO₂) Exchanges in Wheat Using Flux-Tower and Remote Sensing. *J Indian Soc Remote Sens*. 2011;39(3):383–91.
14. Lujikx IT, Neubert REM, Laan Svan der, Meijer HAJ. Continuous measurements of atmospheric oxygen and carbon dioxide on a North sea gas platform. *Atmos Meas Tech Discuss*. 2009;2:1693–724.
15. Sanchez-Canete EP, Serrano-Ortiz P, Kowalski AS, Oyonarte C, Domingo F. Subterranean CO₂ ventilation and its role in the net ecosystem carbon balance of karstic shrubland. *Geophys Res Letters*. 2011;38: L09802, doi: 10.1029/2011GL047077.
16. Sapunova S, Ivanova T, Kostov P, Naydenov Y, Llieva L, Dandolov L. Monitoring and control of atmospheric gas composition in Space plant growth facilities: selection of CO₂ sensors For the svet-3 space greenhouse. *Ecol Eng Environ Protec*. 2008;1:56–64.
17. Zhen J, Chuan-Kuan W, Xing-Chang W. Spatio-temporal variations of CO₂ within the canopy in a temperate deciduous forest, Northeast China. *Chinese J Plant Ecol*. 2009;35:512–22.
18. Hari P, Pumpanen J, Huotari J, et al. High-frequency measurements of productivity of planktonic algae using rugged nondispersive infrared carbon dioxide probes. *Limnol Oceanogr: Methods*. 2008;6:347–54.
19. Vargas R, Baldochi DD, Allen MF, et al. Looking deeper into the soil: biophysical controls and seasonal lags of soil CO₂ production and efflux. *Ecol Appl*. 2010;20(6):1569–82.
20. Vaisala. Vaisala CARBOCAP Carbon Dioxide Probe GMP343. Vaisala Oyj, Finland. 2005. <http://www.vaisala.com>.
21. Mani A. Handbook of Solar Radiation: Data for India. *Allied Publishers Private Limited*. 1980; New Delhi: 500.
22. Agrawal S, Joshi PK, Shukla Y, Roy PS. SPOT VEGETATION multi temporal data for classifying vegetation in south central Asia. *Current Science*. 2003;84:1440–8.



23. Reynolds CA, Jackson TJ, Rawls WJ. Estimated available water content from the FAO soil map of the world, global soil profile databases, pedo-transfer functions. 1999. *Boulder, NOAA National Geophysical Data Center*.
24. Draxler RR, Rolph GD. HySPLIT (Hybrid Single Particle Lagrangian Integrated Trajectory) Model access via NOAA ARL READY website (<http://www.arl.noaa.gov/ready/hysplit4.html>), NOAA Air Resources Laboratory. *Silver Spring, MD*. 2003.
25. Rolph GD. Real-time Environmental Applications and Display System (READY) Website (<http://www.arl.noaa.gov/ready/hysplit4.html>). NOAA Air Resources Laboratory. *Silver Spring, MD*. 2003.
26. Potter CS, Randerson JT, Field CB. Terrestrial ecosystem production: A process model on global satellite and surface data. *Global Biogeochem Cycles*. 1993;7(4):811–41.
27. Field CB, Randerson JT, Malmstrom CM. Global net primary production: Combining ecology and remote sensing. *Remote Sens Environ*. 1995;51(1): 74–88.
28. Thompson MV, Randerson JT, Malstrom CM, Field CB. Change in Net primary production and heterotrophic respiration. How much is necessary to sustain the terrestrial carbon sink? *Global Biogeochem Cycles*. 1996;10(4):711–26.
29. Potter CS, Klooster SA, Myneni RB, Genovesi V, Tan PN, Kumar V. Continental scale comparisons of terrestrial carbon sinks estimated from satellite data and ecosystem modeling 1982–1998. *Global Planet Change*. 2003;39:201–13.
30. Nayak RK, Patel NR, Dadhwal VK. Estimation and analysis of terrestrial net primary productivity over India by remote-sensing-driven terrestrial biosphere model. *Environ Monit Assess*. 2009;170(1–4):195–213.
31. Nayak RK, Patel NR, Dadhwal VK. Inter-annual variability of Net Primary Productivity over India. *Internat J Climat*. 2012;DOI: 10.1002/joc.3414.
32. Bakwin PS, Zhao C, Ussler III W, Tans PP, Quesnel E. Measurements of carbon dioxide on a very tall tower. *Tellus B*. 1995;47(5):535–49.
33. Bakwin PS, Tans PP, White JWC, Andres RJ. Determination the isotopic (13C/12C) discrimination by terrestrial biology from a global net-work of observations. *Global Biogeochem Cycles*. 1998;12:555–62.
34. Spittlehouse DL, Ripley EA. Carbon dioxide concentrations over a native grassland in Saskatchewan. *Tellus*. 1997;29:54–65.
35. Yuesi W, Changke W, Xueqing GUO, Guangren LIU, Yao H. Trend, seasonal and diurnal variations of atmospheric CO₂ in Beijing. *Chinese Science Bulletin*. 2002;47:24.
36. Stull RB. An Introduction to Boundary Layer Meteorology. *Kluwer Acad., Dordrecht, Netherlands*. 1988:667.
37. Ghosh D. Large point sources (lps) emissions from India: Regional and sectoral analysis. *Atmos Environ*. 2002;36:213–24.