

REVIEW

OPEN ACCESS
Full open access to this and
thousands of other papers at
<http://www.la-press.com>.

Bovine Herpes Virus 1 (BHV-1) and Herpes Simplex Virus Type 1 (HSV-1) Promote Survival of Latently Infected Sensory Neurons, in Part by Inhibiting Apoptosis

Clinton Jones

School of Veterinary Medicine and Biomedical Sciences, Nebraska Center for Virology, University of Nebraska, Morrison Life Science Center, Lincoln, NE. Corresponding author email: cjones2@unl.edu

Abstract: α -Herpesvirinae subfamily members, including herpes simplex virus type 1 (HSV-1) and bovine herpes virus 1 (BHV-1), initiate infection in mucosal surfaces. BHV-1 and HSV-1 enter sensory neurons by cell-cell spread where a burst of viral gene expression occurs. When compared to non-neuronal cells, viral gene expression is quickly extinguished in sensory neurons resulting in neuronal survival and latency. The HSV-1 latency associated transcript (LAT), which is abundantly expressed in latently infected neurons, inhibits apoptosis, viral transcription, and productive infection, and directly or indirectly enhances reactivation from latency in small animal models. Three anti-apoptosis genes can be substituted for LAT, which will restore wild type levels of reactivation from latency to a LAT null mutant virus. Two small non-coding RNAs encoded by LAT possess anti-apoptosis functions in transfected cells. The BHV-1 latency related RNA (LR-RNA), like LAT, is abundantly expressed during latency. The LR-RNA encodes a protein (ORF2) and two microRNAs that are expressed in certain latently infected neurons. Wild-type expression of LR gene products is required for stress-induced reactivation from latency in cattle. ORF2 has anti-apoptosis functions and interacts with certain cellular transcription factors that stimulate viral transcription and productive infection. ORF2 is predicted to promote survival of infected neurons by inhibiting apoptosis and sequestering cellular transcription factors which stimulate productive infection. In addition, the LR encoded microRNAs inhibit viral transcription and apoptosis. In summary, the ability of BHV-1 and HSV-1 to interfere with apoptosis and productive infection in sensory neurons is crucial for the life-long latency-reactivation cycle in their respective hosts.

Keywords: alpha-herpesviruses, latency in sensory neurons, apoptosis, non-coding RNAs

Journal of Cell Death 2013:6 1–16

doi: [10.4137/JCD.S10803](https://doi.org/10.4137/JCD.S10803)

This article is available from <http://www.la-press.com>.

© the author(s), publisher and licensee Libertas Academica Ltd.

This is an open access article published under the Creative Commons CC-BY-NC 3.0 license.



α -Herpesvirinae Subfamily Members are Important Pathogens in their Respective Hosts

Herpes simplex virus type 1 (HSV-1) and bovine herpes virus 1 (BHV-1) are both important pathogens in their respective natural hosts and both are α -herpesvirinae subfamily members. For example, HSV-1 is the cause of one of the most frequent and serious viral eye infections in the United States, with over 400,000 affected individuals.¹ Following primary infection of the eye, latency is established in sensory neurons within trigeminal ganglia (TG).^{2,3} HSV-1 reactivates sporadically from TG and the infectious virus can be detected on surfaces of the eye, where it can cause recurrent ocular disease. Reactivation from latency is necessary for recurrent ocular HSV-1 infections.^{4,5} Long-term oral acyclovir treatment only reduces ocular HSV-1 recurrences by 41%.⁶ Herpes simplex virus type 2 (HSV-2) is the cause of the most common sexually transmitted disease, and sporadic recurrent genital lesions occur periodically. Two genital herpes vaccine trials failed^{7,8} indicating there is a need for new and effective therapies that will reduce the incidence of recurrent HSV-1 and HSV-2 disease.

Bovine herpes virus 1 (BHV-1) is an important pathogen of cattle as it induces clinical signs in the upper respiratory tract of cattle and is immunosuppressive. BHV-1 establishes latency in sensory neurons, but periodically reactivates from latency, and thus is widespread in cattle.^{2,9–11} BHV-1 infection inhibits cell-mediated immunity,^{12–15} CD8⁺ T cell recognition of infected cells,^{16–19} and induces apoptosis in CD4⁺ T cells.^{20,21} Two viral regulatory proteins, bICP0 and bICP27, inhibit interferon dependent transcription.^{10,22–25} Infection also erodes mucosal surfaces of the upper respiratory tract, which can allow bacterial pathogens to colonize the lower respiratory tract.^{26–28}

Acute Infection Results in High Levels of Infectious Virus and Apoptosis

Binding and entry of HSV-1 and BHV-1 to mammalian cells are mediated by viral glycoproteins and cellular factors.^{29–31} A cellular receptor (HveA or HVEM) is primarily expressed in activated T cells and belongs to the tumor necrosis factor receptor family.³² Entry of HSV-1 into epithelial and fibroblasts is mediated

by another membrane glycoprotein, HveB or HveC.³³ HveC is an entry mediator for HSV-1 and BHV-1 and is abundantly expressed in neurons. Additionally, soluble HveC blocks viral entry in neuronal-like cell lines.³³ After uncoating, the viral genome enters the nucleus and productive infection is initiated.

HSV-1 and BHV-1 gene expression is tightly regulated in three distinct phases during productive infection of cultured cells: immediate early (IE), early (E), or late (L).³⁴ IE RNA expression does not require protein synthesis and is stimulated by VP16, a viral structural protein.³⁵ E gene expression requires at least one IE protein, and E genes encode nonstructural proteins that stimulate viral DNA replication. L gene expression is maximal after viral DNA replication, requires IE protein production, and L proteins comprise the virion particle. Although a vigorous immune response leads to viral clearance following primary infection, BHV-1 and HSV-1 establish a life-long latent infection in ganglionic sensory neurons, primarily TG, or sacral dorsal root ganglia.^{2,3,9,36} Approximately 40% of sensory neurons appear to harbor viral genomes during latency.^{37–41}

Five HSV-1 IE genes encode ICP0, ICP4, ICP22, ICP27, or ICP47. ICP4^{42–45} and ICP27^{46–48} are required for virus growth in tissue culture. ICP4 represses IE gene expression^{44,49–53} but activates E or L gene expression by interacting with RNA polymerase II transcription factors and specifically binding viral DNA.^{49,54} ICP27 redistributes small nuclear ribonucleoprotein complexes, interferes with splicing of IE transcripts, and promotes E and L poly A site selection.^{55–58} ICP47 prevents transport of antigenic peptides into the endoplasmic reticulum⁵⁹ and inhibits CD8⁺ T cell responses.⁶⁰ ICP22 enhances viral gene expression, in part by modifying RNA polymerase II.⁶¹ ICP0 increases steady-state levels of viral mRNA and stimulates all viral promoters.⁶² ICP0 also binds several cellular proteins: (1) elongation factor 1- α ⁶³; (2) cyclin D3⁶⁴; (3) an ubiquitin-specific protease^{65,66}; and (4) promyelocytic leukemia (PML) protein.^{67–69} Interactions between ICP0 and chromatin-remodeling enzymes activate viral transcription by multiple mechanisms, including sequestering histone deacetylase (HDAC) inhibitors.^{70,71} Secondly, HSV-1 ICP0 interacts with HDAC2⁷² and blocks histone deacetylation to stimulate viral gene expression.^{73,74} ICP0 also alters a complex that inhibits

gene expression (REST/CoREST/HDAC repressor complex).⁷³ Since ICP0 can remove histones from viral chromatin during productive infection,⁷⁵ ICP0 may have similar functions during reactivation from latency. These activities of ICP0 promote virus replication in differentiated cells.⁷⁶ BHV-1 encoded ICP0 (bICP0) has similar functions as ICP0.¹⁰

Viral infection routinely leads to apoptosis in cultured cells.^{77–80} Killing of infected cells by apoptosis *in vivo* can reduce inflammation, alter immune recognition, reduce burst size, and thus prevent virus spread. Premature apoptosis of infected cells limits production of infectious virus and limits viral spread. Members of the α -herpesvirinae subfamily induce apoptosis after infection of cultured cells.^{81–86} HSV-1^{83,84,87–89} and BHV-1⁹⁰ can also inhibit apoptosis in a cell type dependent manner after infection of cultured cells. HSV can induce DNA damage, and consequently apoptosis, even in the absence of productive infection.^{91–95} Two viral proteins, U_S1.5 and U_L13, activate caspase 3 in the absence of other viral proteins, indicating these viral proteins play an important role during virus mediated apoptosis.⁹⁶ Finally, ICP0 is also a trigger for apoptosis in the context of productive infection, in part because it activates viral gene expression.

HSV-1 encodes several proteins (ICP27, U_S3, U_S5, gJ, gD, and LAT) that have anti-apoptosis activity.^{83–85,87,88,97–105} U_S3 is a serine/threonine protein kinase that inhibits cleavage and activation of the

pro-apoptotic Bcl-2 family member, Bad. U_S3 protein expression in cultured cells, in the absence of other viral proteins, inhibits caspase 3 activation, a crucial executioner caspase that commits cells to apoptosis. As expected, U_S3 inhibits the pro-apoptotic activity of U_S1.5 and U_L13 by blocking caspase 3 activation.⁹⁶ These anti-apoptotic genes play an important role in the pathogenic properties of HSV-1.

Viral Genes Expressed During Latency Regulate the Latency-Reactivation Cycle

The HSV-1 latency associated transcript is abundantly expressed during latency and regulates the latency-reactivation cycle

The HSV-1 latency associated transcript (LAT) is abundantly expressed in sensory ganglionic neurons of mice, rabbits, or humans that are latently infected.^{106–114} LAT is predominantly expressed in the nucleus of latently infected neurons suggesting it is a non-protein coding regulatory RNA. LAT is antisense to ICP0 and overlaps ICP0 mRNA sequences (Fig. 1B), suggesting LAT inhibits ICP0 expression by an antisense mechanism. Although the ability of LAT to repress ICP0 expression may be important, LAT sequences that promote spontaneous reactivation in a rabbit ocular model of infection do not overlap ICP0 mRNA sequences.¹¹⁵




Figure 1. Location of genes within the HSV-1 repeats. (Panel A) U_L and U_S denote the unique sequences of the long (L) and short (S) components of the genome. The boxes depict repeat sequences. (Panel B) Transcription map of the repeat region. Location and orientation of LAT,^{111,112} ICP0, $\alpha_134.5$,^{253,254} ORFP,¹⁵⁰ L/STs²⁵⁵ are indicated by solid lines.

Note: Partially mapped transcripts (αX and βX) are denoted by dashed arrows.^{256,257}

Splicing of the 8.5 kb LAT transcript yields a stable 2 kb LAT and an unstable 6.5 kb LAT (Fig. 1B).^{107,111,116} Correct splicing of the 2 kb LAT is necessary for establishment and maintenance of latency.^{117,118} In general, the 2 kb LAT is not capped, is poly A-, appears to be circular, and is a stable intron.^{119,120} A subset of LAT is detected in the cytoplasm^{97,121,122} and is associated with polyribosomes or splicing factors.^{97,123} Small non-coding RNAs can regulate gene expression,^{124,125} promote neuronal differentiation,¹²⁶ or inhibit apoptosis¹²⁷ suggesting LAT is a non-coding regulatory RNA.

A study by Umbach et al¹²⁸ concluded that LAT is a microRNA (miRNA) precursor which encodes four miRNAs, two within LAT promoter sequences (Fig. 2A and B). LAT miR-H6, reduces ICP4 protein

steady state levels but not ICP4 RNA levels. Protein levels, not RNA levels, of ICP0 are inhibited by the LAT miRNA, miR-H2-3p. Within the first 1.5 kb of LAT coding sequences, two small RNAs (sRNAs), LAT sRNA1 and sRNA2, were identified (Fig. 2B). The sRNAs are larger than mature miRNAs (typically 23 nucleotides long) and the sequence of both possess extensive secondary structure. The sRNAs can be detected in TG of mice latently infected with wild-type HSV-1, but not in TG of mice latently infected with a LAT null mutant.^{79,129} LAT sRNA2, but not LAT sRNA1, reduced ICP4 protein levels in transient transfection assays. Both LAT sRNAs inhibit productive infection in mouse neuroblastoma cells, however LAT sRNA1 inhibited productive infection more efficiently than LAT sRNA2.¹³⁰ Collectively, these




Figure 2. Schematic of putative factors encoded within the LAT locus. (Panel A) Schematic of genes within the long repeats that contain the LAT locus. The large arrow indicates the primary LAT transcript. The solid rectangle represents the sTable 2 kb LAT intron. Initiation of LAT transcription is denoted by the arrow at +1 (genomic nucleotide 118801). Several restriction enzyme sites and the relative locations of the ICP0 and ICP34.5 transcripts are shown for reference. The location of the 6 microRNAs (miR-H1-6) that is located within the 8.5 kb LAT¹²⁸ are shown. (Panel B) Partial restriction map of LAT and position of LAT open reading frames (L1-8) within the first 1.5 Kb of strain McKrae LAT coding sequences, which were based on previous studies.¹⁴⁸ The numbering system of the ORFs was consistent with a previous study.¹⁴⁸ Only the ORFs with at least 30 amino acids are shown (the number of amino acids in each ORF is denoted by the numbers in brackets). Open circles denote the position of two LAT small RNAs that are encoded within the first 1.5 kb LAT coding sequences.²⁵⁸ Positions of UOL transcript, AL transcript, and ORFs located on the opposite strand of LAT (AL2 and AL3) are shown. The number of amino acids of AL2 and AL3 are in brackets. Nucleotide positions relative to the start of LAT transcription are not shown in parenthesis.



studies provide evidence that the LAT encoded miRNAs and sRNAs promote latency by interfering with expression of important viral transcriptional regulatory proteins.

The LAT locus encodes additional transcripts (Fig. 2B). More than one transcript, including UOL (Upstream of LAT)¹³¹ are located within LAT promoter sequences. Expression of the UOL transcript or protein does not reduce reactivation from latency in rabbits.¹³² An antisense to LAT (AL) transcript is expressed from the first 1.5 kb of LAT coding sequences and appears to encode a protein.¹³³ Two additional small open reading frames (ORFs) that are antisense to LAT (AL2 and AL3) are present in LAT coding sequences. An AL3 transcript is expressed during productive infection and in TG of mice latently infected with wild-type HSV-1, but not a LAT null mutant virus.¹³⁴ An AL3-specific polyclonal antibody detected a protein in a subset of TG neurons in latently infected mice. A transcript encompassing AL2 has not been detected during productive infection or latency (unpublished data).

LAT null HSV-1 mutants have been examined in various small animal models.^{2,3} Although two studies concluded that LAT does not play a role in latency,^{135,136} most have provided evidence that LAT is important. This discrepancy may be due to the strain of virus or mouse that was used for specific studies. LAT enhances the establishment of latency in mice^{137,138} and in rabbit ocular infection models,¹³⁹ in part by reducing lytic cycle viral gene expression in TG of mice.^{140,141} By enhancing the establishment of latency, LAT would increase the pool of latently infected neurons; thus indirectly increasing the incidence of reactivation from latency.

As a result of stress or other external stimuli, reactivation from latency can occur, resulting in virus shedding (Fig. 4). The McKrae strain of wild-type HSV-1, however not a LAT null mutant, is consistently detected in tears of infected rabbits, due to spontaneous reactivation.^{139,142–145} These same LAT null mutants grow with wild-type efficiency in cultured cells and in acutely infected rabbits. When just the first 1.5 kb of LAT coding sequences (Fig. 2B) is expressed from the HSV-1 genome, wild-type levels of spontaneous reactivation from latency occur in rabbits.¹³⁹ Similar results were observed using another virulent strain of HSV-1 (17 syn+) in a rabbit eye model.^{146,147}

The first 1.5 kb of LAT coding sequences does not overlap ICP0, suggesting that antisense repression of ICP0 expression by LAT is not important for spontaneous reactivation in the rabbit ocular model of infection. The factors encoded by the first 1.5 kb of LAT coding sequences that promote spontaneous reactivation have not been fully characterized.

Although certain studies suggested LAT does not encode a protein,¹⁴⁸ other studies have concluded that a protein encoded within LAT sequences is expressed.^{131,149–154} These proteins were suggested to either substitute for ICP0 functions,^{153,154} interfere with binding of ICP4 to DNA,¹⁵² or their functions were not described. The proposed LAT proteins are mapped downstream of the critical first 1.5 kb of the primary LAT transcript, a region that appears both sufficient and necessary for the wild type spontaneous reactivation phenotype in rabbit models.^{139,155} Within the first 1.5 kb of LAT coding sequences, 8 potential ORFs have been identified in the McKrae strain (Fig. 2B).¹⁴⁸ The L2 ORF (Fig. 2B) appears to be expressed in TG of latently infected mice.¹⁵⁶ Although LAT is not absolutely required for the latency-reactivation cycle in small animal models, its importance may be underestimated using small animal models and measuring latency in terms of weeks or months, not decades.

The BHV-1 latency related RNA is abundantly expressed in sensory neurons and is necessary for reactivation from latency

Latency related (LR) RNA is abundantly expressed in TG neurons of calves that are latently infected.^{111,157} Two different start sites of LR-RNA transcription (Fig. 3A) have been identified suggesting this has functional significance. It is clear that the LR gene encodes more than one product.^{11,36} For example, the LR gene contains two well defined ORFs (ORF2 and ORF1; Fig. 3B) and two reading frames that lack an initiating methionine (RF-B and RF-C). As a result of alternative splicing of polyA+ LR-RNA in TG of infected calves (Fig. 3A),^{158,159} ORF2 can be fused with ORF1 protein coding sequences or RF-B. The ORF2/ORF1 fusion protein stably interacts with the cellular transcription factor C/EBP-alpha.¹⁶⁰ C/EBP-alpha RNA and protein levels increase in TG neurons




Figure 3. Schematic of the BHV-1 LR gene and surrounding genes. (Panel **A**) The start sites for LR transcription during latency and productive infection were previously described.^{159,259} (Panel **B**) Organization of LR ORFs and 3' terminus of bICP0. ORF-1 and ORF-2 are located in the LR gene and have the potential to encode a 40 or 25 kd protein respectively. **Notes:** Reading Frames B (RF-B) and C (RF-C) are open reading frames that lack an initiating Met. The (*) denotes the position of stop codons that are in frame with the respective ORF. The positions of ORF-E and bICP0, which are antisense to LR-RNA, are also shown.

during dexamethasone induced reactivation from latency. Over-expression of C/EBP-alpha enhanced productive infection,¹⁶¹ suggesting that ORF2 sequesters C/EBP-alpha and reduces the efficiency of productive infection during the latency-reactivation cycle.

One day after calves are infected and during latency, splicing of LR-RNA in TG is such that ORF2 is intact,¹⁵⁹ suggesting ORF2 expression is important for the latency-reactivation cycle. ORF2 interacts

with Notch1 and Notch3, components of the Notch signaling pathway.¹⁶² Mammalian Notch receptor family members (Notch1-4) are membrane tethered transcription factors that regulate many developmental and physiological processes.^{163,164} For example, Notch promotes neuronal maintenance, development, and differentiation.¹⁶⁵⁻¹⁶⁷ Notch3¹⁶⁸ and Notch1^{169,170} promote cell survival by activating a protein kinase, (AKT) which inhibits apoptosis. Notch family members can also induce apoptosis,^{163,164} suggesting Notch influences cell survival by cell-type dependent mechanisms. When the Notch receptor is engaged by one of its five transmembrane ligands (Jagged1, Jagged2, Delta-like1, Delta-like3, or Delta-like4), the Notch intracellular domain (ICD) is cleaved by specific proteases, and subsequently translocates to the nucleus. In the nucleus, Notch ICD interacts with members of the CSL family of transcriptional factors, CBF1, Su(H), or Lag1 (also referred to as RBP-J binding proteins) subsequently activating downstream genes. Notch1, but not Notch3, enhances BHV-1 productive infection¹⁶³ and Notch1 activates the BHV-1 immediate-early transcription unit 1 (IEt1) and bICP0 early promoters. Notch1 and Notch3 trans-activated the late glycoprotein C (gC)




Figure 4. Putative steps that occur during the latency-reactivation cycle. **Note:** For details, see text.



promoter. ORF2 interferes with the ability of Notch1 to trans-activate the bICP0 early promoter and Notch1 or Notch3 mediated activation of the gC promoter¹⁶² suggesting this function is important for establishing and/or maintaining latency. Notch3 RNA levels are higher during dexamethasone (DEX) induced reactivation from latency, suggesting Notch family members stimulate productive infection during reactivation from latency. Activation of Notch signaling in post-mitotic-neurons or neuroblastoma cells inhibits neurite sprouting^{165,171–174} and axon repair,¹⁷⁵ which can lead to neuronal degeneration and apoptosis.^{176–178} Conversely, neurite sprouting correlates with regeneration of damaged axons and dendrites.¹⁷⁵ ORF2 promotes neurite sprouting and neuronal differentiation of mouse neuroblastoma cells when Notch1 or Notch3 is over-expressed.¹⁷⁹ Collectively, these studies suggest that ORF2 interactions with Notch family members promote the establishment and maintenance of latency by (1) interfering with viral gene expression necessary for productive infection, (2) supporting a mature neuronal phenotype, and (3) overcoming the deleterious effects of Notch expression during stress-induced reactivation from latency.

Although the results from the LR mutant virus suggested that proteins encoded by the LR gene are necessary for the latency-reactivation cycle, non-protein coding functions within LR-RNA have also been identified. For example, the intact LR gene inhibits the ability of bICP0 to stimulate productive infection in a dose-dependent manner.^{180,181} Insertion of three in-frame stop codons at the amino-terminus of the first ORF within the LR gene (ORF2) inhibited bICP0 repression with similar efficiency as the wild-type LR gene, suggesting expression of a LR protein is not required.¹⁸¹ LR gene products also inhibit mammalian cell growth,^{182,183} and the cell growth inhibitory function of the LR gene maps to a 463-bp fragment that lacks a significant open reading frame.¹⁸² Two miRNAs located upstream of ORF2 are expressed during latency.¹⁸⁴ These miRNAs, or larger sRNAs containing these miRNAs, reduced bICP0 protein levels in transient transfection assays.

A small ORF located within the LR promoter is designated ORF-E (Fig. 3B). ORF-E is antisense to the LR transcript and is downstream of bICP0 coding sequences, but does not overlap bICP0. A transcript

that encompasses ORF-E is expressed during productive infection and in TG of latently infected calves.¹⁸⁵ The LR promoter contains multiple cis-acting motifs, has a neuronal specific binding domain,^{186–188} and contains a long AT-rich motif (40/53 nucleotides are A or T) that may promote ORF-E transcription. When ORF-E protein coding sequences are fused in frame with green fluorescent protein (GFP) sequences, GFP protein expression is detected in the nucleus of mouse or human neuroblastoma cells. In contrast, the ORF-E-GFP fusion protein is detected throughout rabbit skin cells. In transient transfection assays, ORF-E promotes neurite formation in mouse neuroblastoma cells,¹⁸⁹ which may support a mature neuronal phenotype following infection.

LAT and LR Gene Products Inhibit Apoptosis

LAT inhibits apoptosis

LAT expressing plasmids interfere with apoptosis in transiently transfected cells, and LAT expressing viruses inhibit apoptosis in TG of infected mice or rabbits.^{117,190–192} The anti-apoptotic functions of LAT correlate with promoting spontaneous reactivation from latency.^{191,193} In the context of promoting spontaneous reactivation from latency in the rabbit model (model), inhibiting apoptosis is the most important function of LAT as three different anti-apoptosis genes^{129,194–196} restore wild-type levels of spontaneous reactivation from latency to a LAT null mutant. LAT may encode other functions because the LAT null mutants that express cellular anti-apoptosis genes have reduced virulence, in spite of reactivating from latency with wild-type frequency. LAT expressing plasmids, in the absence of other viral genes, inhibit caspase 8- and caspase 9-induced apoptosis,^{193,197} the two major apoptotic pathways in mammals.^{198–200} LAT also inhibits caspase 3 activation.²⁰¹

LAT sRNA1 and sRNA2 cooperate to inhibit cold-shock induced apoptosis in mouse neuroblastoma cells.¹³⁰ Introduction of ATG→TTG mutations in ORFs within the first 1.5 kb of LAT coding sequences impairs the anti-apoptotic functions of LAT,²⁰² suggesting that LAT either encodes a functional protein or alters RNA structure. Two of these ATG→TTG mutations are within LAT sRNA1 and sRNA2, and introducing these mutations into the small RNAs inhibits



their ability to inhibit apoptosis.¹³⁰ At this time, it is not clear how these sRNAs interfere with apoptosis. It will also be important to construct a recombinant virus with these same mutations and test whether the spontaneous reactivation incidence is affected.

LAT also inhibits GrzB induced apoptosis in transient transfection studies.²⁰³ GrzB is released from CD8⁺ T cells as well as other specific lymphocytes; GrzB has features similar to apical caspases, and can induce apoptosis in most cell types.^{204–207} Inhibiting GrzB induced apoptosis may be important for the latency-reactivation cycle because CD8⁺ T lymphocytes control HSV infection in sensory ganglia.^{208,209}

The LR gene encodes more than one product that inhibits apoptosis

A mutant BHV-1 strain with 3 stop codons after the initiating methionine codon of ORF-2 (LR mutant virus) does not express detectable levels of ORF-2¹⁰¹ but expresses reduced levels of ORF1 in cultured cells during productive infection.²¹⁰ The LR mutant virus grows less efficiently in the ocular cavity and TG, but grows almost as efficiently as wild-type BHV-1 in the nasal cavity, and does not reactivate from latency following DEX treatment.^{211,212} The LR mutant virus induces higher levels of apoptosis in TG neurons of infected calves,²¹³ and a LR gene expressing plasmid with the same stop codon mutations does not effectively inhibit apoptosis.^{214,215} ORF2 expression in the absence of other viral genes inhibits apoptosis in transiently transfected cells,^{216,217} suggesting that ORF2 is a dominant function encoded by the LR gene. ORF2, like LAT, can inhibit caspase 8 and caspase 9 mediated apoptosis; however the mechanism by which it inhibits apoptosis is not known.

Two microRNAs encoded within the LR gene (Fig. 3A) interfere with bICP0 protein expression¹⁸⁴ and cold shock induced apoptosis in transfected mouse neuroblastoma cells (Neuro-2A cells).²¹⁸ Since cold shock induced apoptosis in Neuro-2A cells is inhibited by caspase 3 and caspase 9 inhibitors,²¹⁹ the microRNAs must influence these apoptotic signaling pathways. The ability of the microRNAs to stimulate the anti-apoptotic transcription factor NF- α B^{220–223} seems to be important for inhibiting cold-shock induced apoptosis. In summary, these

results provide additional evidence that interfering with apoptosis is crucial for a successful life-long latent infection.

Why is Inhibiting Neuronal Apoptosis Important During the Latency-Reactivation Cycle?

The latency-reactivation cycle has been operationally divided into three distinct steps: establishment, maintenance, and reactivation (Fig. 4). Following acute infection where high levels of infectious virus are produced, virus particles enter sensory neurons. Initial entry of the viral genome into a sensory neuron results in a burst of lytic cycle viral gene expression and infectious viruses are produced. Viral gene expression is then extinguished, with the exception of HSV-1 LAT and BHV-1 LR gene products. Neuronal cell factors,^{2,9} LAT encoded microRNAs plus sRNAs,^{128,130} and LR encoded functions^{162,184} interfere with various aspects of productive infection. During acute infection and establishment of latency, neuronal and satellite cells undergo apoptosis when small animal models are infected with HSV-2²²⁴ or HSV-1.^{225–227} BHV-1 replication and gene expression also occur in TG of acutely infected calves, resulting in apoptosis of neurons and non-neuronal cells.^{192,213,228}

HSV-1 LAT¹⁹² and the LR gene²¹³ enhance neuronal survival during the establishment of latency. The ability of LAT²²⁹ and the LR gene¹⁷⁹ to promote a mature neuronal phenotype and sprout neurites may also promote establishment of latency by stimulating repair of damaged neurons following infection. Successful establishment correlates with an increase in the number of infected neurons that survive and enhances the probability that reactivation from latency occurs.

Maintenance of latency is a phase that lasts for the duration of the host's life and is operationally defined as a period when infectious virus is not readily detected. In general, abundant expression of viral genes required for productive infection does not occur. LAT or LR gene products are abundantly expressed during the maintenance of latency. Expression of LAT correlates with an increase of latently infected neurons during the maintenance of latency,²³⁰ suggesting latently infected sensory neurons are exposed to apoptotic stimuli during the maintenance of latency. It is reasonable to predict that LAT and LR gene products



actively participate in maintaining a latent infection in sensory neurons.

Reactivation from latency is initiated by external stimuli (stress, immunosuppression, or UV light for example), which ultimately must stimulate viral gene expression.^{36,231,232} Abundant viral gene expression can be detected in sensory neurons and infectious virus can be isolated from TG, ocular swabs, and/or nasal swabs. Stress leads to elevated corticosteroid levels, which has rapid effects on neural activity.^{233,234} DEX, a synthetic corticosteroid, induces viral gene expression,²³⁵ stimulates an HSV-1 origin of replication (Ori-L) in neuronal cells,⁵⁵ and alters splicing patterns in the absence of protein synthesis.²³⁶ DEX and other apoptosis stimulators can also stimulate HSV-1 reactivation from latency.^{237,238} BHV-1 reactivation from latency is induced by DEX, in part because it stimulates expression of cellular transcription factors and viral gene expression while repressing expression of LR gene products.^{9,11,36} Prolonged exposure to corticosteroids can also induce immunosuppression, in part by inducing apoptosis in lymphocytes.²³⁹ A subset of neurons that successfully reactivate from latency to produce infectious virus may not survive;^{240–242} however it is not clear if this is the fate for all neurons that produce infectious virus during reactivation from latency. Most latently infected neurons that are exposed to reactivation stimuli re-establish latency and do not produce infectious virus.^{243,244} Given that sensory neurons are terminally differentiated cells, inhibiting apoptosis during the latency-reactivation cycle is crucial for life-long latent infections of α -herpesvirinae subfamily members.

Numerous studies have demonstrated that infiltrating lymphocytes in TG regulate the latency-reactivation cycle. For example, a persistent cell-mediated immune response occurs in TG during latency and CD8⁺ T lymphocytes inhibit reactivation from latency.^{208,209,245–250} Release of granzyme B from CD8⁺ T cells into latently infected neurons helps to inhibit reactivation from latency by cleaving the viral transcriptional trans-activator, ICP4.²⁵¹ Since it is well established that granzyme B activates caspase 3 and the intrinsic pathway of apoptosis,²⁰⁷ the ability of LAT and perhaps LR gene products to inhibit apoptosis is important to overcome the effects of granzyme B. The ability of HSV-1 to inhibit major histocompatibility complex (MHC) class I presentation in sensory

neurons correlates with successful reactivation²⁵² providing further evidence that CD8⁺ T cells monitor latently infected neurons. In conclusion, the ability of HSV-1, HSV-2, and BHV-1 to reactivate from latency is regulated by complex virus-host interactions.

Perspectives

Although genetic and functional studies have demonstrated that LAT and the LR gene regulate the latency-reactivation cycle, there are many unanswered questions. For example, identifying the functions of various transcripts, the small non-coding RNAs, and the ORFs within LAT are crucial to understand the role these various factors play in the latency-reactivation cycle. It would not be surprising to find that one or more of these factors encoded within the LAT locus regulate certain neuronal specific functions that maintain normal functions. It will be difficult to make additional LAT mutant viruses as many of these factors overlap and deletion of these sequences would likely interfere with expression of more than one LAT encoded factor. Consequently, many of these studies will have to be performed in transient transfection assays in primary neurons or neuroblastoma cells. Finally, examining LAT in small animal models in terms of weeks after acute infection may not accurately reflect the latency-reactivation cycle in the context of life-long latency in humans.

With respect to the LR gene, there are no studies that have determined whether ORF1, ORF-E, or the microRNAs play a role in the latency-reactivation cycle. Furthermore, functional analysis of ORF-1 and ORF-E has not been performed. Identifying the cellular proteins that interact with ORF-1 and ORF-E may provide insight into their functions. In summary, the finding that sequences encompassing LAT and the LR gene encode for more than one transcript and/or small non-coding RNAs implies many functions are necessary to successfully regulate the life-long latency-reactivation in the natural host.

Author Contributions

Conceived and designed the experiments: CJ. Analyzed the data: CJ. Wrote the first draft of the manuscript: CJ. Contributed to the writing of the manuscript: CJ. Agree with manuscript results and conclusions: CJ. Jointly developed the structure and arguments for the paper: CJ. Made critical revisions and approved final version: CJ. All authors reviewed and approved of the final manuscript.



Funding

Research in the author's laboratory is supported by a grant from the USDA, NIFA Competitive Grants Program (09-01653), a grant to the Nebraska Center for Virology (1P20RR15635), and the NE Research Initiative.

Competing Interests

Author disclose no potential conflicts of interest.

Disclosures and Ethics

As a requirement of publication the author has provided signed confirmation of compliance with ethical and legal obligations including but not limited to compliance with ICMJE authorship and competing interests guidelines, that the article is neither under consideration for publication nor published elsewhere, of their compliance with legal and ethical guidelines concerning human and animal research participants (if applicable), and that permission has been obtained for reproduction of any copyrighted material. This article was subject to blind, independent, expert peer review. The reviewers reported no competing interests. Provenance: the authors were invited to submit this paper.

References

- Smith RE, McDonald HR, Nesburn AB, Minckler DS. Penetrating keratoplasty: changing indications, 1947 to 1978. *Arch Ophthalmol*. 1980;98(7):1226–9.
- Jones C. Herpes simplex virus type 1 and bovine herpesvirus 1 latency. *Clin Micro Rev*. 2003;16(1):79–95.
- Wagner EK, Bloom DC. Experimental investigation of herpes simplex virus latency. *Clin Microbiol Rev*. 1997;10(3):419–43.
- Nesburn AB. Recurrent herpes simplex infection: pathogenesis and treatment. In: Barraquer J, Binder P, Buxton J, editors. *Symposium on Medical and Surgical Disease of the Cornea*. CV Mosby Company, St. Louis; 1980:48–68.
- Nesburn AB, editor. *Report of the Corneal Disease Panel: Vision Research: A National Plan 1983–1987*, vol. II. CV Mosby Co., St. Louis; 1983.
- Nesburn AB, Ghiasi H, Wechsler SL. Ocular safety and efficacy of an HSV-1 gD vaccine during primary and latent infection. *Invest Ophthalmol Vis Sci*. 1990;31(8):1497–502.
- Cohen J. Painful failure of promising genital herpes vaccine. *Science*. 2010;330(6002):304.
- Stanberry LR, Spruance SL, Cunningham AL, et al. Glycoprotein-D- adjuvant vaccine to prevent genital herpes. *N Engl J Med*. 2002;347(21):1652–61.
- Jones C. Alpha herpesvirus latency: its role in disease and survival of the virus in nature. *Adv Virus Res*. 1998;51:81–133.
- Jones C. Regulation of innate immune responses by bovine herpesvirus 1 and infected cell protein 0. *Viruses*. 2009;1(2):255–75.
- Jones C, Geiser V, Henderson G, et al. Functional analysis of bovine herpesvirus 1 (BHV-1) genes expressed during latency. *Vet Micro*. 2006;113(3–4):199–210.
- Carter JJ, Weinberg AD, Pollard A, Reeves R, Magnuson JA, Magnuson NS. Inhibition of T-lymphocyte mitogenic responses and effects on cell functions by bovine herpesvirus 1. *J Virol*. 1989;63(4):1525–30.
- Griebel P, Ohmann HB, Lawman MJ, Babiuk LA. The interaction between bovine herpesvirus type 1 and activated bovine T lymphocytes. *J Gen Virol*. 1990;71(Pt 2):369–77.
- Griebel P, Qualtiere L, Davis WC, et al. T lymphocyte population dynamics and function following a primary bovine herpesvirus type-1 infection. *Viral Immunol*. 1987;1(4):287–304.
- Griebel P, Qualtiere L, Davis WC, Lawman MJ, Babiuk LA. Bovine peripheral blood leukocyte subpopulation dynamics following a primary bovine herpesvirus-1 infection. *Viral Immunol*. 1987–1988;1(4):267–86.
- Hariharan MJ, Nataraj C, Srikumaran S. Down regulation of murine MHC class I expression by bovine herpesvirus 1. *Viral Immunol*. 1993;6(4):273–84.
- Hinkley S, Hill AB, Srikumaran S. Bovine herpesvirus-1 infection affects the peptide transport activity in bovine cells. *Virus Res*. 1998;53(1):91–6.
- Koppers-Lalic EA, Reits EAJ, Rensing ME, et al. Varicelloviruses avoid T cell recognition by UL49.5-mediated inactivation of the transporter associated with antigen processing. *Proc Natl Acad Sci U S A*. 2005;102(14):5144–9.
- Nataraj C, Eidmann S, Hariharan MJ, Sur JH, Perry GA, Srikumaran S. Bovine herpesvirus 1 downregulates the expression of bovine MHC class I molecules. *Viral Immunol*. 1997;10(1):21–34.
- Eskra L, Splitter GA. Bovine herpesvirus-1 infects activated CD4+ lymphocytes. *J Gen Virol*. 1997;78(Pt 9):2159–66.
- Winkler MT, Doster A, Jones C. Bovine herpesvirus 1 can infect CD4(+) T lymphocytes and induce programmed cell death during acute infection of cattle. *J Virol*. 1999;73:8657–68.
- da Silva LF, Sinani D, Jones C. The ICP27 protein encoded by bovine herpesvirus type 1 (bICP27) interferes with promoter activity of the bovine genes encoding beta interferon 1 (IFN- β 1) and IFN- β 3. *Virus Res*. 2012;169(1):162–8.
- Henderson G, Zhang Y, Jones C. The bovine herpesvirus 1 gene encoding infected cell protein 0 (bICP0) can inhibit interferon-dependent transcription in the absence of other viral genes. *J Gen Virol*. 2005;86(Pt 10):2697–702.
- Saira K, Zhou Y, Jones C. The infected cell protein 0 encoded by bovine herpesvirus 1 (bICP0) induces degradation of interferon response factor 3 (IRF3), and consequently inhibits beta interferon promoter activity. *J Virol*. 2007;81(7):3077–86.
- Saira K, Jones C. The infected cell protein 0 encoded by bovine herpesvirus 1 (bICP0) associates with interferon regulatory factor 7 (IRF7), and consequently inhibits beta interferon promoter activity. *J Virol*. 2009;83(8):3977–81.
- Highlander SK. Molecular genetic analysis of virulence in Mannheimia (Pasteurella) haemolytica. *Front Biosci*. 2001;D1128–50.
- Highlander SK, Fedorova ND, Dusek DM, Panciera R, Alvarez LE, Renehart C. Inactivation of Pasteurella (Mannheimia) haemolytica leukotoxin causes partial attenuation of virulence in a calf challenge model. *Infect Immun*. 2000;68(7):3916–22.
- Zecchinon L, Fett T, Desmecht D. How Mannheimia haemolytica defeats host defense through a kiss of death mechanism. *Vet Res*. 2005;36(2):133–56.
- Spear PG. Herpes simplex virus: receptors and ligands for cell entry. *Cell Microbiol*. 2004;6(5):401–10.
- Spear PG, Manoj S, Yoon M, Jogger CR, Zago A, Myscofski D. Different receptors binding to distinct interfaces on herpes simplex virus gD can trigger events leading to cell fusion and viral entry. *Virology*. 2006;344(1):17–24.
- Spear PG, Longnecker R. Herpesvirus entry: an update. *J Virol*. 2003;77(19):10179–85.
- Montgomery RI, Warner MS, Lum BJ, Spear PG. Herpes simplex virus-1 entry into cells mediated by a novel member of the TNF/NGF receptor family. *Cell*. 1996;87(3):427–36.
- Geraghty RJ, Krummenacher C, Cohen GH, Eisenberg RJ, Spear PG. Entry of alpha herpesviruses mediated by poliovirus receptor-related protein 1 and poliovirus receptor. *Science*. 1998;280(5369):1618–20.
- Honess RW, Roizman B. Regulation of herpes virus macromolecular synthesis: Cascade regulation of three groups of viral proteins. *J Virol*. 1974;14(1):8–19.
- O'Hare P. The virion transactivator of herpes simplex virus. *Seminars Virol*. 1993;4:145–55.



36. Jones C, da Silva LF, Sinani D. Regulation of the latency-reactivation cycle by products encoded by the bovine herpesvirus 1 (BHV-1) latency-related gene. *J Neurovirol.* 2011;17(6):535–45.
37. Maggioncalda J, Mehta A, Su YH, Fraser NW, Block TM. Correlation between herpes simplex virus type 1 rate of reactivation from latent infection and the number of infected neurons in trigeminal ganglia. *Virology.* 1996;225(1):72–81.
38. Mehta A, Maggioncalda J, Bagasra O, et al. In situ DNA PCR and RNA hybridization detection of herpes simplex virus sequences in trigeminal ganglia of latently infected mice. *Virology.* 1995;206(1):633–40.
39. Ramakrishnan R, Fink DJ, Jiang G, Desai P, Glorioso JC, Levine M. Competitive quantitative PCR analysis of herpes simplex virus type 1 DNA and latency-associated transcript RNA in latently infected cells of the rat brain. *J Virol.* Mar 1994;68(3):1864–73.
40. Ramakrishnan R, Levine M, Fink DJ. PCR-based analysis of herpes simplex virus type 1 latency in the rat trigeminal ganglion established with a ribonucleotide reductase-deficient mutant. *J Virol.* 1994;68:7083–91.
41. Sawtell NM. Comprehensive quantification of herpes simplex virus latency at the single-cell level. *J Virol.* 1997;71(7):5423–31.
42. Carozza MJ, DeLuca NA. Interaction of the viral activator protein ICP4 with TFIID through TAF250. *Mol Cell Biol.* 1996;16(6):3085–93.
43. DeLuca NA, McCarthy AM, Schaffer PA. Isolation and characterization of deletion mutants of herpes simplex virus type 1 in the gene encoding immediate-early regulatory protein ICP4. *J Virol.* 1985;56(2):558–70.
44. DeLuca NA, Schaffer PA. Activation of immediate-early, early, and late promoters by temperature-sensitive and wild-type forms of herpes simplex virus type 1 protein ICP4. *Mol Cell Biol.* 1985;5(8):1997–208.
45. Dixon RA, Schaffer PA. Fine-structure mapping and functional analysis of temperature-sensitive mutants in the gene encoding the herpes simplex virus type 1 immediate early protein VP175. *J Virol.* 1980;36(1):189–203.
46. McCarthy AM, McMahan L, Schaffer PA. Herpes simplex virus type 1 ICP27 deletion mutants exhibit altered patterns of transcription and are DNA deficient. *J Virol.* 1989;63(1):18–27.
47. McMahan L, Schaffer PA. The repressing and enhancing functions of the herpes simplex virus regulatory protein ICP27 map to C-terminal regions and are required to modulate viral gene expression very early in infection. *J Virol.* 1990;64(7):3471–85.
48. Sacks WR, Greene CC, Aschman DP, Schaffer PA. Herpes simplex virus type 1 ICP27 is an essential regulatory protein. *J Virol.* 1985;55(3):796–805.
49. Gu B, DeLuca N. Requirements for activation of the herpes simplex virus glycoprotein C promoter in vitro by the viral regulatory protein ICP4. *J Virol.* 1994;68(12):7953–65.
50. Gu B, Rivera-Gonzalez R, Smith CA, DeLuca NA. Herpes simplex virus infected cell polypeptide 4 preferentially represses Sp1-activated over basal transcription from its own promoter. *Proc Natl Acad Sci U S A.* 1993; 90(20):9528–32.
51. Michael N, Roizman B. Repression of the herpes simplex virus 1 alpha 4 gene by its gene product occurs within the context of the viral genome and is associated with all three identified cognate sites. *Proc Natl Acad Sci U S A.* 1993;90(6):2286–90.
52. O'Hare P, Hayward GS. Three trans-acting regulatory proteins of herpes simplex virus modulate immediate-early gene expression in a pathway involving positive and negative feedback regulation. *J Virol.* 1985;56(3):723–33.
53. Roberts MS, Boundy A, O'Hare P, Pizzorno MC, Ciuffo DM, Hayward GS. Direct correlation between a negative autoregulatory response element at the cap site of the herpes simplex virus type 1 IE175 (alpha 4) promoter and a specific binding site for the IE175 (ICP4) protein. *J Virol.* 1988;62(11):4307–20.
54. Smith CA, Bates P, Rivera-Gonzalez R, Gu B, DeLuca NA. ICP4, the major transcriptional regulatory protein of herpes simplex virus type 1, forms a tripartite complex with TATA-binding protein and TFIIB. *J Virol.* 1993;67(8): 4676–87.
55. Hardwicke MA, Schaffer PA. Differential effects of nerve growth factor and dexamethasone on herpes simplex virus type 1 oriL- and oriS-dependent DNA replication in PC12 cells. *J Virol.* 1997;71(5):3580–7.
56. Hardy WR, Sandri-Goldin RM. Herpes simplex virus inhibits host cell splicing, and regulatory protein ICP27 is required for this effect. *J Virol.* 1994;68(12):7790–9.
57. Sandri-Goldin RM, Hibbard MK, Hardwicke MA. The C-terminal repressor region of herpes simplex virus type 1 ICP27 is required for the redistribution of small nuclear ribonucleoprotein particles and splicing factor SC35; however, these alterations are not sufficient to inhibit host cell splicing. *J Virol.* 1995;69(10):6063–76.
58. Sandri-Goldin RM, Mendoza GE. A herpesvirus regulatory protein appears to act post-transcriptionally by affecting mRNA processing. *Genes Dev.* 1992;6(5):848–63.
59. Hill A, Jugovic P, York I, et al. Herpes simplex virus turns off the TAP to evade host immunity. *Nature.* 1995;375(6530):411–5.
60. Goldsmith K, Chen W, Johnson DC, Hendricks RL. Infected cell protein (ICP)47 enhances herpes simplex virus neurovirulence by blocking the CD8+ T cell response. *J Exp Med.* 1998;187(3):341–8.
61. Rice SA, Long MC, Lam V, Schaffer PA, Spencer CA. Herpes simplex virus immediate-early protein ICP22 is required for viral modification of host RNA polymerase II and establishment of the normal viral transcription program. *J Virol.* 1995;69(9):5550–9.
62. Jordan R, Schaffer PA. Activation of gene expression by herpes simplex virus type 1 ICP0 occurs at the level of mRNA synthesis. *J Virol.* 1997;71(9): 6850–62.
63. Kawaguchi Y, Bruni R, Roizman B. Interaction of herpes simplex virus 1 alpha regulatory protein ICP0 with elongation factor 1 delta: ICP0 affects translational machinery. *J Virol.* 1997;71(2):1019–24.
64. Kawaguchi Y, Van Sant C, Roizman B. Herpes simplex virus 1 alpha regulatory protein ICP0 interacts with and stabilizes the cell cycle regulator cyclin D3. *J Virol.* 1997;71(10):7328–36.
65. Meredith M, Orr A, Elliott M, Everett R. Separation of sequence requirements for HSV-1 Vmw110 multimerisation and interaction with a 135-kDa cellular protein. *Virology.* 1995;209(1):174–87.
66. Meredith M, Orr A, Everett R. Herpes simplex virus type 1 immediate-early protein Vmw110 binds strongly and specifically to a 135-kDa cellular protein. *Virology.* 1994;200(2):457–69.
67. Everett R, O'Hare P, O'Rourke D, Barlow P, Orr A. Point mutations in the herpes simplex virus type 1 Vmw110 RING finger helix affect activation of gene expression, viral growth, and interaction with PML-containing nuclear structures. *J Virol.* 1995;69(11):7339–44.
68. Everett RD, Lomonte P, Sternsdorf T, van Driel R, Orr A. Cell cycle regulation of PML modification and ND10 composition. *J Cell Sci.* 1999; 112(Pt 24):4581–8.
69. Everett RD, Meredith M, Orr A, Cross A, Kathoria M, Parkinson J. A novel ubiquitin-specific protease is dynamically associated with the PML nuclear domain and binds to a herpesvirus regulatory protein. *EMBO J.* 1997;16(7):1519–30.
70. Hobbs WE, DeLuca NA. Perturbation of cell cycle progression and cellular gene expression as a function of herpes simplex virus ICP0. *J Virol.* 1999;73(10):8245–55.
71. Poon AP, Liang Y, Roizman B. Herpes simplex virus 1 gene expression is accelerated by inhibitors of histone deacetylases in rabbit skin cells infected with a mutant carrying a cDNA copy of the infected cell protein no. *J Virol.* 2003;77(23):12671–8.
72. Lomonte P, Seigneurin-Berny D, Everett RD, Khochbin S, Epstein AL. Presented at the 26th International Herpesvirus Workshop; 2001.
73. Gu H, Liang Y, Mandel G, Roizman B. Components of the REST/CoREST/histone deacetylase repressor complex are disrupted, modified, and translocated in HSV-1-infected cells. *Proc Nat Acad Sci U S A.* 2005;102(21):7571–6.
74. Poon APW, Gu H, Roizman B. ICP0 and the Us3 protein kinase of herpes simplex virus 1 independently block histone deacetylation to enable gene expression. *Proc Nat Acad Sci U S A.* 2006;103(26):9993–8.
75. Cliffe AR, Knipe DM. Herpes simplex virus ICP0 promotes both histone removal and acetylation on viral dna during lytic infection. *J Virol.* 2008;82(24):12030–8.
76. Cai W, Schaffer PA. A cellular function can enhance gene expression and plating efficiency of a mutant defective in the gene for ICP0, a transactivating protein of herpes simplex virus type 1. *J Virol.* 1991;65(8):4078–90.
77. Hardwick JM. Viral interference with apoptosis. *Semin Cell Dev Biol.* 1998;9(3):339–49.



78. Razvi ES, Welsh RM. Apoptosis in viral infections. *Adv Virus Res.* 1995; 45:1–60.
79. Shen Y, Shenk TE. Viruses and apoptosis. *Curr Opin Genet Dev.* 1995;5: 105–11.
80. Teodoro JG, Branton PE. Regulation of apoptosis by viral gene products. *J Virol.* 1997;71(3):1739–46.
81. Aiamkitsumrit B, Zhang X, Block TM, Norton P, Fraser NW, Su YH. Herpes simplex virus type 1 ICP4 deletion mutant virus d120 infection failed to induce apoptosis in nerve growth factor-differentiated PC12 cells. *J Neurovirol.* 2007;13(4):305–14.
82. Devireddy LR, Jones C. Activation of caspases and p53 by bovine herpesvirus 1 infection results in programmed cell death and efficient virus release. *J Virol.* 1999;73(5):3778–88.
83. Galvan V, Brandimarti R, Roizman B. Herpes simplex virus 1 blocks caspase-3-independent and caspase-dependent pathways to cell death. *J Virol.* 1999;73(4):3219–26.
84. Galvan V, Roizman B. Herpes simplex virus 1 induces and blocks apoptosis at multiple steps during infection and protects cells from exogenous inducers in a cell-type-dependent manner. *Proc Natl Acad Sci U S A.* 1998;95(7):3931–6.
85. Pradhan P, Nguyen ML. Early passage neonatal and adult keratinocytes are sensitive to apoptosis induced by infection with an ICP27-null mutant of herpes simplex virus 1. *Apoptosis.* 2013;18(2):160–70.
86. Sadzot-Delvaux C, Thonard P, Schoonbroodt S, Piette J, Rentier B. Varicella-zoster virus induces apoptosis in cell culture. *J Gen Virol.* 1995; 76(Pt 11):2875–9.
87. Asano S, Honda T, Goshima F, et al. US3 protein kinase of herpes simplex virus type 2 plays a role in protecting corneal epithelial cells from apoptosis in infected mice. *J Gen Virol.* 1999;80(Pt 1):51–6.
88. Aubert M, Blaho JA. The herpes simplex virus type 1 regulatory protein ICP27 is required for the prevention of apoptosis in infected human cells. *J Virol.* 1999;73(4):2803–13.
89. Leopardi R, Roizman B. The herpes simplex virus major regulatory protein ICP4 blocks apoptosis induced by the virus or by hyperthermia. *Proc Natl Acad Sci U S A.* 1996;93(18):9583–7.
90. Geiser V, Rose S, Jones C. The bovine herpes virus 1 bICP0 protein regulates toxicity in a cell type dependent fashion. *Molec Path.* 2008;44:459–66.
91. Chenet-Monte C, Mohammad F, Celluzzi CM, Schaffer PA, Farber FE. Herpes simplex virus gene products involved in the induction of chromosomal aberrations. *Virus Res.* 1986;6(3):245–60.
92. Ellison SA, Hampar B. Chromosomal aberrations induced by an animal virus. *Nature.* 1961;192:145–7.
93. Heilbronn R, zur Hausen H. A subset of herpes simplex virus replication genes induces DNA amplification within the host cell genome. *J Virol.* 1989;63(9):3683–92.
94. Pilon L, Langelier Y, Royat A. Herpes simplex virus type 2 mutagenesis: characterization of mutants induced at the hprt locus of nonpermissive XC cells. *Mol Cell Biol.* 1986;6(8):2977–83.
95. Soengas MS, Alarcon RM, Yoshida H, et al. Apaf-1 and caspase-9 in p53-dependent apoptosis and tumor inhibition. *Science.* 1999;284(5411): 156–9.
96. Hagglund R, Munger J, Poon AP, Roizman B. U(S)3 protein kinase of herpes simplex virus 1 blocks caspase 3 activation induced by the products of U(S)1.5 and U(L)13 genes and modulates expression of transduced U(S)1.5 open reading frame in a cell type-specific manner. *J Virol.* 2002;76(2):743–54.
97. Ahmed M, Fraser NW. Herpes simplex virus type 1 2-kilobase latency-associated transcript intron associates with ribosomal proteins and splicing factors. *J Virol.* 2001;75(24):12070–80.
98. Blaho JA, Aubert M. Modulation of apoptosis during herpes simplex virus infection in human cells. *Microbes Infect.* 2001;3(10):859–66.
99. Jerome KR, Chen Z, Lang R, et al. HSV and Glycoprotein J inhibit caspase activation and apoptosis induced by Granzyme B or Fas. *J Immunol.* 2001;167(7):3928–35.
100. Jerome KR, Fox R, Chen Z, Sears AE, Lee H, Corey L. Herpes simplex virus inhibits apoptosis through the action of two genes, Us5 and Us3. *J Virol.* 1999;73(11):8950–7.
101. Jiang Y, Inman M, Zhang Y, Posadas NA, Jones C. A mutation in the latency related gene of bovine herpesvirus 1 (BHV-1) inhibits protein expression of a protein from open reading frame 2 (ORF-2) and an adjacent reading frame during productive infection. *J Virol.* 2004;78(6):3184–9.
102. Kather A, Raftery MJ, Devi-Rao G, et al. Herpes simplex virus type 1 (HSV-1)-induced apoptosis in human dendritic cells as a result of downregulation of cellular FLICE-inhibitory protein and reduced expression of HSV-1 antiapoptotic latency-associated transcript sequences. *J Virol.* 2010;84(2):1034–46.
103. Munger J, Chee AV, Roizman B. The U(S)3 protein kinase blocks apoptosis induced by the d120 mutant of herpes simplex virus 1 at a premitochondrial stage. *J Virol.* 2001;75(12):5491–7.
104. Munger J, Roizman B. The US3 protein kinase of herpes simplex virus 1 mediates the posttranslational modification of BAD and prevents BAD-induced programmed cell death in the absence of other viral proteins. *Proc Natl Acad Sci U S A.* 2001;98(18):10410–5.
105. Nguyen ML, Blaho JA. Cellular players in the herpes simplex virus dependent apoptosis balancing act. *Viruses.* 2009;1(3):965–78.
106. Croen KD, Ostrove JM, Dragovic LJ, Smialek JE, Straus SE. Latent herpes simplex virus in human trigeminal ganglia. Detection of an immediate early gene “anti-sense” transcript by in situ hybridization. *N Engl J Med.* 1987;317(23):1427–32.
107. Deatly AM, Spivack JG, Lavi E, O’Boyle DR 2nd, Fraser NW. Latent herpes simplex virus type 1 transcripts in peripheral and central nervous system tissues of mice map to similar regions of the viral genome. *J Virol.* 1988;62(3):749–56.
108. Deatly AM, Spivack JG, Lavi E, Fraser NW. RNA from an immediate early region of the type 1 herpes simplex virus genome is present in the trigeminal ganglia of latently infected mice. *Proc Natl Acad Sci U S A.* 1987;84(10):3204–8.
109. Krause PR, Croen KD, Straus SE, Ostrove JM. Detection and preliminary characterization of herpes simplex virus type 1 transcripts in latently infected human trigeminal ganglia. *J Virol.* 1988;62(12):4819–23.
110. Mitchell WJ, Lirette RP, Fraser NW. Mapping of low abundance latency-associated RNA in the trigeminal ganglia of mice latently infected with herpes simplex virus type 1. *J Gen Virol.* 1990;71(Pt 1):125–32.
111. Rock DL, Nesburn AB, Ghiasi H, et al. Detection of latency-related viral RNAs in trigeminal ganglia of rabbits latently infected with herpes simplex virus type 1. *J Virol.* 1987;61(12):3820–6.
112. Stevens JG, Wagner EK, Devi-Rao GB, Cook ML, Feldman LT. RNA complementary to a herpesvirus alpha gene mRNA is prominent in latently infected neurons. *Science.* 1987;235(4792):1056–9.
113. Wagner EK, Devi-Rao G, Feldman LT, et al. Physical characterization of the herpes simplex virus latency-associated transcript in neurons. *J Virol.* 1988;62(4):1194–202.
114. Wagner EK, Flanagan WM, Devi-Rao G, et al. The herpes simplex virus latency-associated transcript is spliced during the latent phase of infection. *J Virol.* 1988;62(12):4577–85.
115. Perng GC, Ghiasi H, Slanina SM, Nesburn AB, Wechsler SL. The spontaneous reactivation function of the herpes simplex virus type 1 LAT gene resides completely within the first 1.5 kilobases of the 8.3-kilobase primary transcript. *J Virol.* 1996;70(2):976–84.
116. Zwaagstra JC, Ghiasi H, Slanina SM, et al. Activity of herpes simplex virus type 1 latency-associated transcript (LAT) promoter in neuron-derived cells: evidence for neuron specificity and for a large LAT transcript. *J Virol.* 1990;64(10):5019–28.
117. Kang W, Mukerjee R, Fraser NF. Establishment and maintenance of HSV latent infection is mediated through correct splicing of the LAT primary transcript. *Virology.* 2003;312(1):233–44.
118. Mador N, Panet A, Latchman D, Steiner I. Expression and splicing of the latency-associated transcripts of herpes simplex virus type 1 in neuronal and non-neuronal cell lines. *J Biochem (Tokyo).* 1995;117(6):1288–97.
119. Farrell MJ, Dobson AT, Feldman LT. Herpes simplex virus latency-associated transcript is a stable intron. *Proc Natl Acad Sci U S A.* 1991;88: 790–4.
120. Krummenacher C, Zabolotny JM, Fraser NW. Selection of a nonconsensus branch point is influenced by an RNA stem-loop structure and is important to confer stability to the herpes simplex virus 2-kilobase latency-associated transcript. *J Virol.* 1997;71(8):5849–60.



121. Nicosia M, Zabolotny JM, Lirette RP, Fraser NW. The HSV-1 2-kb latency-associated transcript is found in the cytoplasm comigrating with ribosomal subunits during productive infection. *Virology*. 1994;204(2):717–28.
122. Thomas DL, Lock M, Zabolotny JM, Mohan BR, Fraser NW. The 2-kilobase intron of the herpes simplex virus type 1 latency-associated transcript has a half-life of approximately 24 hours in SY5Y and COS-1 cells. *J Virol*. 2002;76(2):532–40.
123. Goldenberg D, Mador N, Ball MJ, Panet A, Steiner I. The abundant latency-associated transcripts of herpes simplex virus type 1 are bound to polyribosomes in cultured neuronal cells and during latent infection in mouse trigeminal ganglia. *J Virol*. 1997;71(4):2897–904.
124. Dykxhoorn DM, Novina CD, Sharp PA. Killing the messenger: short RNAs that silence gene expression. *Nat Rev Mol Cell Biol*. 2003;4(6):457–67.
125. Hannon GJ. RNA interference. *Nature*. 2002;418(6894):244–51.
126. Kuwabera T, Hsieh J, Nakashima K, Taira K, Gage FH. A small modulatory dsRNA specified the fate of adult neural stem cells. *Cell*. 2004;116(6):779–93.
127. Xu P, Guo M, Hay BA. MicroRNAs and the regulation of cell death. *Trends Genet*. 2004;20(12):617–24.
128. Umbach JL, Kramer MF, Jurak I, Karnowski HW, Coen DM, Cullen BR. MicroRNAs expressed by herpes simplex virus 1 during latent infection regulate viral mRNAs. *Nature*. 2008;454(7205):780–5.
129. Perng GC, Maguen B, Jing L, et al. A gene capable of blocking apoptosis can substitute for the herpes simplex virus type 1 latency-associated transcript gene and restore wild-type reactivation levels. *J Virol*. 2002;76(3):1224–35.
130. Shen W, Sa e Silva MSE, Jaber T, et al. Two small RNAs encoded within the first 1.5 kb of the herpes simplex virus type 1 (HSV-1) latency-associated transcript (LAT) can inhibit productive infection, and cooperate to inhibit apoptosis. *J Virol*. 2009;83(18):9131–9.
131. Naito J, Mukerjee R, Mott KR, et al. Identification of a protein encoded in the herpes simplex virus type 1 latency associated transcript promoter region. *Virus Research*. 2005;108(1–2):101–10.
132. Chan D, Cohen J, Naito J, et al. A mutant deleted for most of the herpes simplex virus type 1 (HSV-1) UOL gene does not affect the spontaneous reactivation phenotype in rabbits. *J Neurovirology*. 2006;12(1):5–16.
133. Perng GC, Maguen B, Jing L, et al. A novel herpes simplex virus type 1 (HSV-1) transcript (AL-RNA) antisense to the 5' end of LAT (latency associated transcript) produces a protein in infected rabbits. *J Virol*. 2002;76(16):8003–10.
134. Jaber T, Henderson G, Li S, et al. Identification of a novel herpes simplex virus type 1 (HSV-1) transcript and protein (AL3) expressed during latency. *J Gen Virol*. 2009;90(Pt 10):2342–52.
135. Block TM, Spivack JG, Steiner I, et al. A herpes simplex virus type 1 latency-associated transcript mutant reactivates with normal kinetics from latent infection. *J Virol*. 1990;64(7):3417–26.
136. Ho DY, Mocarski ES. Beta-galactosidase as a marker in the peripheral and neural tissues of the herpes simplex virus-infected mouse. *Virology*. 1988;167(1):279–83.
137. Sawtell NM, Thompson RL. Herpes simplex virus type 1 latency-associated transcription unit promotes anatomical site-dependent establishment and reactivation from latency. *J Virol*. 1992;66(4):2157–69.
138. Thompson RL, Sawtell NM. The herpes simplex virus type 1 latency-associated transcript gene regulates the establishment of latency. *J Virol*. 1997;71(7):5432–40.
139. Perng GC, Ghiasi H, Slanina SM, Nesburn AB, Wechsler SL. The spontaneous reactivation function of the herpes simplex virus type 1 LAT gene resides completely within the first 1.5 kilobases of the 8.3 kilobase primary transcript. *J Virol*. 1996;70(2):976–84.
140. Chen SH, Kramer MF, Schaffer PA, Coen DM. A viral function represses accumulation of transcripts from productive-cycle genes in mouse ganglia latently infected with herpes simplex virus. *J Virol*. 1997;71(8):5878–84.
141. Garber DA, Schaffer PA, Knipe DM. A LAT-associated function reduces productive-cycle gene expression during acute infection of murine sensory neurons with herpes simplex virus type 1. *J Virol*. 1997;71(8):5885–93.
142. Perng GC, Dunkel EC, Geary PA, et al. The latency-associated transcript gene of herpes simplex virus type 1 (HSV-1) is required for efficient in vivo spontaneous reactivation of HSV-1 from latency. *J Virol*. 1994;68(12):8045–55.
143. Perng GC, Slanina S, Ghiasi H, Nesburn AB, Wechsler SL. The effect of latency-associated transcript on the herpes simplex virus type 1 latency-reactivation phenotype is mouse strain-dependent. *J Gen Virol*. 2002;82(Pt 5):1117–22.
144. Perng GC, Slanina AM, Yukht A, et al. A herpes simplex virus type 1 latency-associated transcript mutant with increased virulence and reduced spontaneous reactivation. *J Virol*. 1999;73(2):920–9.
145. Perng GC, Slanina SM, Ghiasi H, Nesburn AB, Wechsler SL. A 371-nucleotide region between the herpes simplex virus type 1 (HSV-1) LAT promoter and the 2-kilobase LAT is not essential for efficient spontaneous reactivation of latent HSV-1. *J Virol*. 1996;70(3):2014–8.
146. Hill JM, Sedarati F, Javier RT, Wagner EK, Stevens JG. Herpes simplex virus latent phase transcription facilitates in vivo reactivation. *Virology*. 1990;174(1):117–25.
147. Trousdale MD, Steiner I, Spivack JG, et al. In vivo and in vitro reactivation impairment of a herpes simplex virus type 1 latency-associated transcript variant in a rabbit eye model. *J Virol*. 1991;65(12):6989–93.
148. Drolet BS, Perng GC, Cohen J, et al. The region of the herpes simplex virus type 1 LAT gene involved in spontaneous reactivation does not encode a functional protein. *Virology*. 1998;242(1):221–32.
149. Doerig C, Pizer LI, Wilcox CL. An antigen encoded by the latency-associated transcript in neuronal cell cultures latently infected with herpes simplex virus type 1. *J Virol*. 1991;65(5):2724–7.
150. Lagunoff M, Roizman B. Expression of a herpes simplex virus 1 open reading frame antisense to the gamma(1)34.5 gene and transcribed by an RNA 3' coterminal with the unspliced latency-associated transcript. *J Virol*. 1994;68(9):6021–8.
151. Lock M, Miller C, Fraser NW. Analysis of protein expression from within the region encoding the 2.0-kilobase latency-associated transcript of herpes simplex virus type 1. *J Virol*. 2001;75(7):3413–26.
152. Randall G, Lagunoff M, Roizman B. The product of ORF O located within the domain of herpes simplex virus 1 genome transcribed during latent infection binds to and inhibits in vitro binding of infected cell protein 4 to its cognate DNA site. *Proc Natl Acad Sci U S A*. 1997;94:10379–84.
153. Thomas SK, Gough G, Latchman DS, Coffin RS. Herpes simplex virus latency-associated transcript encodes a protein which greatly enhances virus growth, can compensate for deficiencies in immediate-early gene expression, and is likely to function during reactivation from virus latency. *J Virol*. 1999;73(8):6618–25.
154. Thomas SK, Lilley CE, Latchman DS, Coffin RS. A Protein Encoded by the Herpes Simplex Virus (HSV) Type 1 2-Kilobase Latency-Associated Transcript Is Phosphorylated, Localized to the Nucleus, and Overcomes the Repression of Expression from Exogenous Promoters When Inserted into the Quiescent HSV Genome. *J Virol*. 2002;76(8):4056–67.
155. Perng GC, Esmail D, Slanina S, et al. Three herpes simplex virus type 1 latency-associated transcript mutants with distinct and asymmetric effects on virulence in mice compared with rabbits. *J Virol*. 2001;75(19):9018–28.
156. Henderson G, Jaber T, Carpenter D, Wechsler SL, Jones C. Identification of herpes simplex virus type 1 (HSV-1) proteins encoded within the first 1.5 kb of the latency-associated transcript (LAT). *J Neurovirology*. 2009;15(5–6):479–88.
157. Rock DL, Beam SL, Mayfield JE. Mapping bovine herpesvirus type 1 latency-related RNA in trigeminal ganglia of latently infected rabbits. *J Virol*. 1987;61(12):3827–31.
158. Devireddy L, Zhang Y, Jones C. Cloning and initial characterization of an alternatively spliced transcript encoded by the bovine herpes virus 1 latency related (LR) gene. *J Neurovirology*. 2003;9(6):612–22.
159. Devireddy LR, Jones C. Alternative splicing of the latency-related transcript of bovine herpesvirus 1 yields RNAs containing unique open reading frames. *J Virol*. 1998;72(9):7294–301.
160. Meyer F, Perez S, Geiser V, Sintek M, Inman M, Jones C. A protein encoded by the bovine herpes virus 1 (BHV-1) latency related gene interacts with specific cellular regulatory proteins, including the CCAAT enhancer binding protein alpha (C/EBP- α). *J Virol*. 2007;81(1):59–67.



161. Meyer F, Jones C. C/EBP-alpha cooperates with bTIF to activate the bovine herpesvirus 1 immediate early transcription unit 1 promoter. *J Neuro Virology*. 2008;2:1–8.
162. Workman A, Sinani D, Pittayakhajonwut D, Jones C. A Protein (ORF2) Encoded by the Latency Related Gene of Bovine Herpesvirus 1 Interacts with Notch1 and Notch3. *J Virol*. 2011;85:2536–46.
163. Bray SJ. Notch signalling: a simple pathway becomes complex. *Nat Rev Mol Cell Biol*. 2006;7(9):678–89.
164. Ehebauer M, Penelope P, Arias AM. Notch, a universal arbiter of cell fate decisions. *Science*. 2006;314(5804):1414–5.
165. Berezovska O, McLean P, Knowles R, et al. Notch1 inhibits neurite outgrowth in postmitotic primary neurons. *Neuroscience*. 1999;93(2):433–9.
166. Cornell R, Eisen JS. Notch in the pathway: the roles of Notch signalling in neural crest development. *Semin Cell Dev Biol*. 2005;16(6):663–72.
167. Justice NJ, Jan YN. Variations on the Notch pathway in neural development. *Curr Op Neurobiol*. 2002;12(1):64–70.
168. Wang T, Holt CM, Xu C, et al. Notch 3 activation modulates growth behavior and cross talks to Wnt/TCF signalling pathway. *Cell Signal*. 2007;19(12):2458–67.
169. Nair P, Somasundaram K, Krishna S. Activated Notch1 inhibits p53-induced apoptosis and sustains transformation by human papilloma virus type 16 E6 and E7 oncogenes through a PI3K-PKB/Akt-dependent pathway. *J Virol*. 2003;77(12):7106–12.
170. Sade H, Krishna S, Sarin A. The anti-apoptotic effect of Notch-1 requires p56lck-dependent, AKT/PKB-mediated signaling in T cells. *J Biol Chem*. 2004;279(4):2937–44.
171. Franklin JL, Berechid BE, Cutting FB, et al. Autonomous and non-autonomous regulation of mammalian neurite development by Notch1 and Delta1. *Curr Biol*. 1999;9(24):1448–57.
172. Levy OA, Lah JJ, Levy AI. Notch signaling inhibits PC12 cell neurite outgrowth via RBP-J-dependent and -independent mechanisms. *Dev Neurosci*. 2002;24:79–88.
173. Levy OA, Lah JJ, Levey AI. Notch signaling inhibits PC12 cell neurite outgrowth via RBP-J-dependent and -independent mechanisms. *Dev Neurosci*. 2002;24:79–88.
174. Sestan N, Artavanis-Tsakonas S, Rakic P. Contact-dependent inhibition of cortical neurite growth mediated by Notch signaling. *Science*. 1999;286(5440):741–6.
175. Coleman MP, Freeman MR. Wallerian degeneration, wld(S), and nmnat. *Annu Rev Neurosci*. 2010;33:245–67.
176. Raff MC, Whitmore AV, Finn JT. Axonal self-destruction and neurodegeneration. *Science*. 2002;296(5569):868–71.
177. Yang Y, Klein R, Tian X, Cheng HT, Kopan R, Shen J. Notch activation induces apoptosis in neural progenitor cells through a p53-dependent pathway. *Developmental Biol*. 2004;269(1):81–04.
178. El Bejjani R, Hammarlund M. Notch signaling inhibits axon regeneration. *Neuron*. 2012;73(2):268–78.
179. Sinani D, Frizzo da Silva L, Jones C. A bovine herpesvirus 1 protein expressed in latently infected neurons (ORF2) promotes neurite sprouting in the presence of activated Notch1 or Notch3. *J Virol*. 2013;87(2):1183–92.
180. Bratanich AC, Hanson ND, Jones C. The latency-related gene of bovine herpesvirus 1 inhibits the activity of immediate-early transcription unit 1. *Virology*. 1992;191(2):988–91.
181. Geiser V, Inman M, Zhang Y, Jones C. The latency related (LR) gene of bovine herpes virus 1 (BHV-1) can inhibit the ability of bICP0 to activate productive infection. *J Gen Virol*. 2002;83:2965–71.
182. Geiser V, Jones C. The latency related gene encoded by bovine herpesvirus 1 encodes a small regulatory RNA that inhibits cell growth. *J Neurovirol*. 2005;11:563–70.
183. Schang LM, Hossain A, Jones C. The latency-related gene of bovine herpesvirus 1 encodes a product which inhibits cell cycle progression. *J Virol*. 1996;70(6):3807–14.
184. Jaber T, Workman A, Jones C. Small noncoding RNAs encoded within the bovine herpesvirus 1 latency-related gene can reduce steady-state levels of infected cell protein 0 (bICP0). *J Virol*. 2010;84(13):6297–307.
185. Inman M, Zhou J, Webb H, Jones C. Identification of a novel transcript containing a small open reading frame that is expressed during latency, and is antisense to the latency related gene of bovine herpes virus 1 (BHV-1). *J Virol*. 2004;78(10):5438–47.
186. Bratanich AC, Jones CJ. Localization of cis-acting sequences in the latency-related promoter of bovine herpesvirus 1 which are regulated by neuronal cell type factors and immediate-early genes. *J Virol*. 1992;66(10):6099–106.
187. Delhon G, Jones C. Identification of DNA sequences in the latency related promoter of bovine herpes virus type 1 which are bound by neuronal specific factors. *Virus Res*. 1997;51(1):93–103.
188. Jones C, Delhon G, Bratanich A, Kutish G, Rock D. Analysis of the transcriptional promoter which regulates the latency-related transcript of bovine herpesvirus 1. *J Virol*. 1990;64(3):1164–70.
189. Perez S, Meyer F, Henderson G, et al. A protein encoded by the bovine herpesvirus 1 ORF E gene induces neurite-like morphological changes in mouse neuroblastoma cells and is expressed in trigeminal ganglionic neurons. *J Neuro Virology*. 2007;13(2):139–49.
190. Ahmed M, Lock M, Miller CG, Fraser NW. Regions of the herpes simplex virus type 1 latency-associated transcript that protect cells from apoptosis in vitro and protect neuronal cells in vivo. *J Virol*. 2002;76(2):717–29.
191. Inman M, Perng GC, Henderson G, et al. Region of herpes simplex virus type 1 latency-associated transcript sufficient for wild-type spontaneous reactivation promotes cell survival in tissue culture. *J Virol*. 2001;75(8):3636–46.
192. Perng GC, Jones C, Ciacci-Zanella J, et al. Virus-induced neuronal apoptosis blocked by the herpes simplex virus latency-associated transcript (LAT). *Science*. 2000;287(5457):1500–3.
193. Jin L, Peng W, Perng GC, Nesburn AB, Jones C, Wechsler SL. Identification of herpes simplex virus type 1 (HSV-1) latency associated transcript (LAT) sequences that both inhibit apoptosis and enhance the spontaneous reactivation phenotype. *J Virol*. 2003;77(11):6556–61.
194. Jin L, Carpenter D, Moerdyk-Schauwecker M, et al. Cellular FLIP can substitute for the herpes simplex virus type 1 LAT gene to support a wild type virus reactivation phenotype in mice. *J Neurovirology*. 2008;14(5):389–400.
195. Jin L, Perng GC, Nesburn AB, Jones C, Wechsler SL. The baculovirus inhibitor of apoptosis gene (cIAP) can restore reactivation of latency to a herpes simplex virus type 1 that does not express the latency associated transcript. *J Virol*. 2005;79(19):12286–95.
196. Mott K, Osorio N, Jin L, et al. The bovine herpesvirus 1 LR ORF2 is crucial for this gene's ability to restore the high reactivation phenotype to a Herpes simplex virus-1 LAT null mutant. *J Gen Virol*. 2003;84:2975–85.
197. Henderson G, Peng W, Jin L, et al. Regulation of caspase 8- and caspase 9-induced apoptosis by the herpes simplex virus latency-associated transcript. *J Neurovirology*. 2002;8 Suppl 2:103–11.
198. Kruegger A, Baumann S, Krammer PH, Kirchhoff S. FLICE-Inhibitory proteins: regulators of death receptor-mediated apoptosis. *Molec Cell Biol*. 2001;21(24):8247–54.
199. Schmitz I, Kirchhoff S, Krammer PH. Regulation of death receptor-mediated apoptosis pathways. *Int J Biochem Cell Biol*. 2000;32(11–12):1123–36.
200. Wang X. The expanding role of mitochondria in apoptosis. *Gen Devel*. 2001;15(22):2922–33.
201. Carpenter D, Hsiang C, Jin L, et al. Stable cell lines expressing high levels of the herpes simplex virus type 1 LAT are refractory to caspase 3 activation and DNA laddering following cold shock induced apoptosis. *Virology*. 2007;369(1):12–8.
202. Carpenter D, Henderson G, Hsiang C, et al. Introducing point mutations into the ATGs of the putative open reading frame of the HSV-1 gene encoding the latency associated transcript (LAT) reduces its anti-apoptotic activity. *Microb Pathog*. 2008;44(2):98–102.
203. Jiang X, Chentoufi A, Hsiang C, et al. The herpes simplex virus type 1 latency associated transcript (LAT) can protect cells from Granzyme B induced apoptosis and CD8 T-cell killing. *J Virol*. 2011;85(5):2325–32.



204. Alimonti JB, Shi L, Baijal PK, Greenberg AH. Granzyme B induces BID-mediated cytochrome c release and mitochondrial permeability transition. *J Biol Chem.* 2001;276(10):6974–82.
205. Barry M, Heibein JA, Pinkoski MJ, et al. Granzyme B short-circuits the need for caspase 8 activity during granule-mediated cytotoxic T-lymphocyte killing by directly cleaving Bid. *Mol Cell Biol.* 2000;20(11):3781–94.
206. Pinkoski MJ, Waterhouse NJ, Heibein JA, et al. Granzyme B-mediated apoptosis proceeds predominantly through a Bcl-2-inhibitable mitochondrial pathway. *J Biol Chem.* 2001;276(15):12060–7.
207. Yang X, Stennicke HR, Wang B, et al. Granzyme B mimics apical caspases. Description of a unified pathway for trans-activation of executioner caspase-3 and -7. *J Biol Chem.* 1998;273:34278–83.
208. Nash AA, Jayasuriya A, Phelan J, Cobbold SP, Waldmann H, Prospero T. Different roles for L3T4+ and Lyt 2+ T cell subsets in the control of an acute herpes simplex virus infection of the skin and nervous system. *J Gen Virol.* 1987;68(Pt 3):825–33.
209. Simmons A, Tschärke D, Speck P. The role of immune mechanisms in control of herpes simplex virus infection of the peripheral nervous system. *Curr Top Microbiol Immunol.* 1992;179:31–56.
210. Meyer F, Perez S, Jiang Y, Zhou Y, Henderson G, Jones C. Identification of a novel protein encoded by the latency-related gene of bovine herpesvirus 1. *J Neuro Virology.* 2007;13(6):569–78.
211. Inman M, Lovato L, Doster A, Jones C. A mutation in the latency related gene of bovine herpesvirus 1 interferes with the latency-reactivation cycle of latency in calves. *J Virol.* 2002;76(13):6771–9.
212. Inman M, Lovato L, Doster A, Jones C. A mutation in the latency-related gene of bovine herpesvirus 1 leads to impaired ocular shedding in acutely infected calves. *J Virol.* 2001;75(18):8507–15.
213. Lovato L, Inman M, Henderson G, Doster A, Jones C. Infection of cattle with a bovine herpesvirus 1 (BHV-1) strain that contains a mutation in the latency related gene leads to increased apoptosis in trigeminal ganglia during the transition from acute infection to latency. *J Virol.* 2003;77(8):4848–57.
214. Ciacci-Zanella J, Stone M, Henderson G, Jones C. The latency-related gene of bovine herpesvirus 1 inhibits programmed cell death. *J Virol.* 1999;73(12):9734–40.
215. Henderson G, Perng GC, Nesburn A, Wechsler S, Jones C. The latency related gene of bovine herpesvirus 1 can suppress caspase 3 and caspase 9 during productive infection. *J Neurovirol.* 2004;10(1):64–70.
216. Shen W, Jones C. Open reading frame 2 encoded by the latency related gene of bovine herpesvirus 1 has anti-apoptosis activity in transiently transfected neuroblastoma cells. *J Virol.* 2008;82:10940–5.
217. Sinani D, Jones C. Localization of sequences in a protein encoded by the latency related gene of bovine herpesvirus 1 (ORF2) that inhibits apoptosis and interferes with Notch1 mediated trans-activation of the bICP0 promoter. *J Virol.* 2011;85(23):12124–33.
218. Silva LF, Jones C. Two microRNAs encoded within the BHV-1 latency related (LR) gene promote cell survival by interacting with RIG-I and stimulating nuclear factor-kappa B (NF-kB) dependent transcription and beta-interferon signaling pathways. *J Virol.* 2012;86(3):1670–82.
219. Shen W, Jones C. Open reading frame 2, encoded by the latency-related gene of bovine herpesvirus 1, has antiapoptotic activity in transiently transfected neuroblastoma cells. *J Virol.* 2008;82(21):10940–5.
220. Antwerp DJ, Martin SJ, Kafri T, Green DR, Verma IM. Suppression of TNF-alpha-induced apoptosis by NF-kB. *Science.* 1996;274(5288):787–9.
221. Foehr ED, Lin X, O'Mahony A, Gelezianus R, Bradshaw RA, Greene WC. NF-kB signaling promotes both cell survival and neurite process formation in nerver growth factor-stimulated PC12 cells. *J Neuroscience.* 2000;20(20):7556–63.
222. Mattson MP, Meffert MK. Roles for NF-kB in nerve cell survival, plasticity, and disease. *Cell Death Differentiation.* 2006;13(5):852–60.
223. Yang CH, Murti A, Pfeffer SR, Basu L, Kim JG, Pfeffer LM. IFNalpha/beta promotes cell survival by activating NF-kappa B. *Proc Natl Acad Sci U S A.* 2000;97(25):13631–6.
224. Ozaki N, Sugiura Y, Yamamoto M, Yokoya S, Wanaka A, Nishiyama Y. Apoptosis induced in the spinal cord and dorsal root ganglion by infection of herpes simplex virus type 2 in the mouse. *Neurosci Lett.* 1997;228:99–102.
225. Knotts FB, Cook ML, Stevens JG. Pathogenesis of herpetic encephalitis in mice after ophthalmic inoculation. *J Infect Dis.* 1974;130(1):16–27.
226. Kramer MF, Chen SH, Knipe DM, Coen DM. Accumulation of viral transcripts and DNA during establishment of latency by herpes simplex virus. *J Virol.* 1998;72(2):1177–85.
227. Speck PG, Simmons A. Divergent molecular pathways of productive and latent infection with a virulent strain of herpes simplex virus type 1. *J Virol.* 1991;65(8):4001–5.
228. Schang L, Jones C. Analysis of bovine herpesvirus 1 transcripts during a primary infection of trigeminal ganglia of cattle. *J Virol.* 1997;71(9):6786–95.
229. Li S, Carpenter D, Hsiang C, Wechsler SL, Jones C. The herpes simplex virus type 1 latency-associated transcript (LAT) locus inhibits apoptosis and promotes neurite sprouting in neuroblastoma cells following serum starvation by maintaining active AKT (protein kinase B). *J Gen Virol.* 2010;91:858–66.
230. Branco FJ, Fraser NW. Herpes simplex virus type 1 latency-associated transcript expression protects trigeminal ganglion neurons from apoptosis. *J Virol.* 1993;79(14):9019–25.
231. Jones C, Chowdhury S. A review of the biology of bovine herpesvirus type 1 (BHV-1), its role as a cofactor in the bovine respiratory disease complex, and development of improved vaccines. *Adv Anim Health.* 2007;8(2):187–205.
232. Perng GC, Jones C. Towards an understanding of the Herpes Simplex Virus Type 1 latency-reactivation cycle. *Interdiscip Perspect Infect Dis.* 2010;2010:262415.
233. Joels M, de Kloet ER. Control of neuronal excitability by corticosteroid hormones. *Trends Neurosci.* 1992;15(1):25–30.
234. McEwen BS. Non-genomic and genomic effects of steroids on neural activity. *Trends Pharmacol Sci.* 1991;12(4):141–7.
235. Halford WP, Gebhardt BM, Carr DJ. Mechanisms of herpes simplex virus type 1 reactivation. *J Virol.* 1996;70(8):5051–60.
236. Collett JW, Steele RE. Alternative splicing of a neural-specific Src mRNA (Src+) is a rapid and protein synthesis-independent response to neural induction in *Xenopus laevis*. *Dev Biol.* 1993;158(2):487–95.
237. Du T, Zhou G, Roizman B. Induction of apoptosis accelerates reactivation from latent HSV-1 in ganglionic organ cultures and replication in cell cultures. *Proc Natl Acad Sci U S A.* 2012;109(36):14616–21.
238. Hunsperger EA, Wilcox CL. Caspase-3-dependent reactivation of latent herpes simplex virus type 1 in sensory neuronal cultures. *J Neurovirology.* 2003;9(3):390–8.
239. Dieken ES, Miesfeld RL. Transcriptional transactivation functions localized to the glucocorticoid receptor N terminus are necessary for steroid induction of lymphocyte apoptosis. *Mol Cell Biol.* 1992;12(2):589–97.
240. Shimeld C, Easty DL, Hill TJ. Reactivation of herpes simplex virus type 1 in the mouse trigeminal ganglion: an in vivo study of virus antigen and cytokines. *J Virol.* 1999;73(3):1767–73.
241. Shimeld C, Hill TJ, Blyth WA, Easty DL. Reactivation of latent infection and induction of recurrent herpetic eye disease in mice. *J Gen Virol.* 1990;71(2):397–404.
242. Shimeld C, Whiteland JL, Williams NA, Easty DL, Hill TJ. Reactivation of herpes simplex virus type 1 in the mouse trigeminal ganglion: an in vivo study of virus antigen and immune cell infiltration. *J Gen Virol.* 1996;77(3):2583–90.
243. Feldman LT, Ellison AR, Voytek CC, Yang L, Krause P, Margolis TP. Spontaneous molecular reactivation of herpes simplex virus type 1 latency in mice. *Proc Natl Acad Sci U S A.* 2002;99(2):978–83.
244. Rock D, Lokensgard J, Lewis T, Kutish G. Characterization of dexamethasone-induced reactivation of latent bovine herpesvirus 1. *J Virol.* 1992;66(4):2484–90.
245. Khanna KM, Bonneau RH, Kinchington PR, Hendricks RL. Herpes simplex virus-specific memory CD8+ T cells are selectively activated and retained in latently infected sensory ganglia. *Immunity.* 2003;18(5):593–603.



246. Liu T, Khanna KM, Carriere BN, Hendricks RL. Gamma interferon can prevent herpes simplex virus type 1 reactivation from latency in sensory neurons. *J Virol.* 2001;75(22):11178–84.
247. Liu T, Khanna KM, Chen X, Fink DJ, Hendricks RL. CD8(+) T cells can block herpes simplex virus type 1 (HSV-1) reactivation from latency in sensory neurons. *J Exp Med.* 2000;191(9):1459–66.
248. Liu T, Tang Q, Hendricks RL. Inflammatory infiltration of the trigeminal ganglion after herpes simplex virus type 1 corneal infection. *J Virol.* 1996;70(1):264–71.
249. Prbhakaran K, Sheridan BS, Kinchington PR, et al. Sensory neurons regulate the effector functions of CD8+ T cells in controlling HSV-1 latency ex vivo. *Immunity.* 2005;23(5):515–23.
250. Simmons A, Tscharke DC. Anti-CD8 impairs clearance of herpes simplex virus from the nervous system: implications for the fate of virally infected neurons. *J Exp Med.* 1992;175(5):1337–44.
251. Knickelbein JE, Khanna KM, Yee MB, Baty CJ, Kinchington PR, Hendricks RL. Noncytotoxic lytic granule-mediated CD8+ T cell inhibition of HSV-1 reactivation from neuronal latency. *Science.* 2008;322(5899):268–72.
252. Orr MT, Mathis MA, Lagunoff M, Sacks JA, Wilson CB. CD8 T cell control of HSV reactivation from latency is abrogated by viral inhibition of MHC Class I. *Cell Host Microbe.* 2007;2(3):172–80.
253. Chou J, Kern ER, Whitley RJ, Roizman B. Mapping of herpes simplex virus-1 neurovirulence to gamma 134.5, a gene nonessential for growth in culture. *Science.* 1990;250(4985):1262–6.
254. Chou J, Roizman B. The herpes simplex virus 1 gene for ICP34.5, which maps in inverted repeats, is conserved in several limited-passage isolates but not in strain 17 syn+. *J Virol.* 1990;64(3):1014–20.
255. Yeh L, Schaffer PA. A novel class of transcripts expressed with late kinetics in the absence of ICP4 spans the junction between the long and short segments of the herpes simplex virus type 1 genome. *J Virol.* 1993;67(12):7373–82.
256. Bohenzky RA, Lagunoff M, Roizman B, Wagner EK, Silverstein S. Two overlapping transcription units which extend across the L-S junction of herpes simplex virus type 1. *J Virol.* 1995;69(5):2889–97.
257. Bohenzky RA, Papavassiliou AG, Gelman IH, Silverstein S. Identification of a promoter mapping within the reiterated sequences that flank the herpes simplex virus type 1 UL region. *J Virol.* 1993;67(2):632–42.
258. Peng W, Vitvitskaia O, Carpenter D, Wechsler SL, Jones C. Identification of two small RNAs within the first 1.5-kb of the herpes simplex virus type 1 (HSV-1) encoded latency-associated transcript (LAT). *J Neuro Virology.* 2008;14(1):41–52.
259. Hossain A, Schang LM, Jones C. Identification of gene products encoded by the latency-related gene of bovine herpesvirus 1. *J Virol.* 1995;69(9):5345–52.